



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>C12N 9/00</b>	<b>A1</b>	(11) International Publication Number: <b>WO 99/61590</b> (43) International Publication Date: 2 December 1999 (02.12.99)
<p>(21) International Application Number: PCT/US99/11780</p> <p>(22) International Filing Date: 28 May 1999 (28.05.99)</p> <p>(30) Priority Data: 60/087,220 29 May 1998 (29.05.98) US</p> <p>(71) Applicant (for all designated States except US): THE SCRIPPS RESEARCH INSTITUTE [US/US]; 10550 North Torrey Pines Road, La Jolla, CA 92037 (US).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): CHERESH, David, A. [US/US]; 327 Via Andalusia, Encinitas, CA 92024 (US). ELICEIRI, Brian [US/US]; 935 Laguna #5, Carlsbad, CA 92008 (US). SCHWARTZBERG, Pamela, L. [US/US]; 49/4A38, National Humane Genome Research Institute, National Institute of Health, Bethesda, MD 20892-4472 (US).</p> <p>(74) Agents: CEPURITIS, Talivaldis et al.; Olson &amp; Hierl, Ltd., 36th floor, 20 North Wacker Drive, Chicago, IL 60606 (US).</p>		<p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p><b>Published</b> <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>
<p>(54) Title: METHODS AND COMPOSITIONS USEFUL FOR MODULATION OF ANGIOGENESIS USING TYROSINE KINASE SRC</p>		
<p>(57) Abstract</p> <p>The present invention describes methods for modulating angiogenesis in tissues using Src protein, modified Src protein, and nucleic acids encoding for such. Particularly the invention describes methods for inhibiting angiogenesis using an inactive Src protein, or nucleic acids encoding therefor, or for potentiating angiogenesis using an active Src protein, or nucleic acids encoding therefor. The invention also describes the use of gene delivery systems for providing nucleic acids encoding for the Src protein, or modified forms thereof.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## METHODS AND COMPOSITIONS USEFUL FOR MODULATION OF ANGIOGENESIS USING TYROSINE KINASE SRC

### Reference to Related Application

5        This application claims the benefit of U.S. Provisional Application Serial No. 60/087,220 filed May 29, 1998, which is incorporated by reference, as are all references cited herein.

### Technical Field

10        The present invention relates generally to the field of medicine, and relates specifically to methods and compositions for modulating angiogenesis of tissues using the protein tyrosine kinase Src, variants of Src, and nucleic acids encoding them.

### Background

15        Angiogenesis is a process of tissue vascularization that involves the growth of new developing blood vessels into a tissue, and is also referred to as neo-vascularization. The process is mediated by the infiltration of endothelial cells and smooth muscle cells. The process is believed to proceed in any one of three ways: the vessels can sprout from pre-existing vessels, de-novo development of vessels can arise from precursor cells (vasculogenesis), or existing small vessels can  
20        enlarge in diameter. Blood et al., Bioch. Biophys. Acta, 1032:89-118 (1990).

      Angiogenesis is an important process in neonatal growth, but is also important in wound healing and in the pathogenesis of a large variety of clinical diseases including tissue inflammation, arthritis, tumor growth, diabetic retinopathy, macular degeneration by neovascularization of the retina and like conditions.  
25        These clinical manifestations associated with angiogenesis are referred to as angiogenic diseases. Folkman et al., Science, 235:442-447 (1987). Angiogenesis is generally absent in adult or mature tissues, although it does occur in wound healing and in the corpus luteum growth cycle. See, for example, Moses et al., Science, 248:1408-1410 (1990).

30        It has been proposed that inhibition of angiogenesis would be a useful therapy for restricting tumor growth. Inhibition of angiogenesis has been proposed by (1) inhibition of release of "angiogenic molecules" such as bFGF (basic fibroblast growth factor), (2) neutralization of angiogenic molecules, such as by use of anti-

$\beta$ bFGF antibodies, (3) use of inhibitors of vitronectin receptor  $\alpha_v\beta_3$ , and (4) inhibition of endothelial cell response to angiogenic stimuli. This latter strategy has received attention, and Folkman et al., Cancer Biology, 3:89-96 (1992), have described several endothelial cell response inhibitors, including collagenase inhibitor, basement membrane turnover inhibitors, angiostatic steroids, fungal-derived angiogenesis inhibitors, platelet factor 4, thrombospondin, arthritis drugs such as D-penicillamine and gold thiomalate, vitamin D<sub>3</sub> analogs, alpha-interferon, and the like that might be used to inhibit angiogenesis. For additional proposed inhibitors of angiogenesis, see Blood et al., Bioch. Biophys. Acta., 1032:89-118 (1990), Moses et al., Science, 248:1408-1410 (1990), Ingber et al., Lab. Invest., 59:44-51 (1988), and United States Patent Nos. 5,092,885, 5,112,946, 5,192,744, 5,202,352, 5,753,230 and 5,766,591. None of the inhibitors of angiogenesis described in the foregoing references involve the Src proteins.

For angiogenesis to occur, endothelial cells must first degrade and cross the blood vessel basement membrane in a similar manner used by tumor cells during invasion and metastasis formation.

It has been previously reported that angiogenesis depends on the interaction between vascular integrins and extracellular matrix proteins. Brooks et al., Science, 264:569-571 (1994). Furthermore, it was reported that programmed cell death (apoptosis) of angiogenic vascular cells is initiated by the interaction, which would be inhibited by certain antagonists of the vascular integrin  $\alpha_v\beta_3$ . Brooks et al., Cell, 79:1157-1164 (1994). More recently, it has been reported that the binding of matrix metalloproteinase-2 (MMP-2) to vitronectin receptor ( $\alpha_v\beta_5$ ) can be inhibited using  $\alpha_v\beta_5$  antagonists, and thereby inhibit the enzymatic function of the proteinase. Brooks et al., Cell, 85:683-693 (1996).

#### Summary of the Invention

The present invention is directed to modulation of angiogenesis in tissues by tyrosine kinase Src, also referred to generically herein as Src.

Compositions and methods for modulating angiogenesis in a tissue associated with a disease condition are contemplated. A composition comprising an angiogenesis-modulating amount of a Src protein is administered to tissue to be treated for a disease condition that responds to modulation of angiogenesis. The composition providing the Src protein can contain purified protein, biologically

active protein fragments, recombinantly produced Src protein or protein fragments or fusion proteins thereof, or gene/nucleic acid expression vectors for expressing a Src protein.

Where the Src protein is inactivated or inhibited, the modulation is an inhibition of angiogenesis. Where the Src protein is active or activated, the modulation is a potentiation of angiogenesis.

The tissue to be treated can be any tissue in which modulation of angiogenesis is desirable. For angiogenesis inhibition, it is useful to treat diseased tissue where deleterious neovascularization is occurring. Exemplary tissues include inflamed tissue, solid tumors, metastases, tissues undergoing restenosis, and the like tissues.

For potentiation, it is useful to treat patients with ischemic limbs in which there is poor circulation in the limbs from diabetic or other conditions. Patients with chronic wounds that do not heal and therefore could benefit from the increase in vascular cell proliferation and neovascularization can be treated as well.

Particularly preferred is the use of Src protein containing a modified amino acid sequence as described herein. Several particularly useful modified Src proteins and the expression thereof are described herein.

The present invention also encompasses a pharmaceutical composition for stimulating angiogenesis in a target mammalian tissue comprising a viral or non-viral gene transfer vector containing a nucleic acid and a pharmaceutically acceptable carrier or excipient; said nucleic acid having a nucleic acid segment encoding for a src protein, said src protein having any amino acid residue at codon 527 except tyrosine, serine or threonine.

Also envisioned is a pharmaceutical composition for inhibiting angiogenesis in a target mammalian tissue comprising a viral or non-viral gene transfer vector containing a nucleic acid and a pharmaceutically acceptable carrier or excipient; said nucleic acid having a nucleic acid segment encoding for a src protein having no kinase activity.

#### Brief Description of the Drawings

In the drawings forming a portion of this disclosure:

FIG. 1 is a cDNA sequence of chicken c-Src which is the complete coding sequence with the introns deleted as first described by Takeya et al., Cell, 32:881-890 (1983). The sequence is accessible through GenBank Accession Number

J00844. The sequence contains 1759 nucleotides with the protein coding portion beginning and ending at the respective nucleotide positions 112 and 1713.

FIG. 2 is the encoded amino acid residue sequence of chicken c-Src of the coding sequence shown in FIG. 1.

5        FIG. 3 is a cDNA sequence of human c-Src which as first described by Braeuninger et al., Proc. Natl. Acad. Sci., USA, 88:10411-10415 (1991). The sequence is accessible through GenBank Accession Number X59932 X71157. The sequence contains 2187 nucleotides with the protein coding portion beginning and ending at the respective nucleotide positions 134 and 1486.

10       FIG. 4 is the encoded amino acid residue sequence of human c-Src of the coding sequence shown in FIG. 3.

FIG. 5 illustrates the activation of endogenous Src by bFGF or VEGF as described in Example 4. The top portion of the figure indicates the results of an in vitro kinase assay with the fold activation of endogenous c-Src by either bFGF and VEGF. The bottom portion of the figure is the kinase assay blot probed with an anti-Src antibody as a loading control for equivalent Src and IgG content.

15       FIG. 6 illustrates the effect of retrovirus-mediated gene expression of c-Src A on angiogenesis in the chick chorioallantoic membrane (CAM) as described in Example 4. Nine-day-old chick CAMs were exposed to RCAS-Src A (active mutated c-Src) or control RCAS-GFP (Green Fluorescent Protein; a fluorescent indicator protein) retroviruses or buffer for 72 h. The level of angiogenesis was quantified as shown in FIG. 6A with representative photomicrographs (4x) in FIG. 6B corresponding to each treatment taken with a stereomicroscope.

20       FIG. 7 illustrates the retroviral expression of c-Src A in activating vascular MAP kinase phosphorylation. FIG. 7A shows tissue extracts of 10 day-old chick CAMs that had been exposed to VEGF or PMA for 30 minutes or infected with c-Src A retrovirus for 48 hours. NT stands for no treatment. Src was immunoprecipitated from equivalent amounts of total protein extract and subjected to an in vitro immune complex kinase assay using a FAK-GST fusion protein as a substrate, electrophoresed and transferred to nitrocellulose. Aliquots of the above whole tissue lysates were also measured for endogenous ERK phosphorylation by immunoblotting with an anti-phospho-ERK antibody. FIG. 7B shows 10 day old CAMs that were infected with either mock RCAS or RCAS containing SRC A.

25      

30

After two days, CAMs were dissected, cryopreserved in OCT and sectioned at 4  $\mu$ m. Sections were immunostained with an anti-phosphorylated ERK antibody (New England Biolabs), washed and detected with a goat anti-rabbit FITC-conjugated secondary antibody. Florescent images were captured on a cooled-CCD camera (Princeton Inst.)

FIG. 8 illustrates the selective requirement for Src activity during VEGF, but not bFGF-induced angiogenesis. Nine day old chick CAMs were exposed to RCAS-Src 251 or control RCAS-GFP retroviruses or buffer for 20 hours and then incubated for an additional 72 hours in the presence or absence of bFGF or VEGF. The level of angiogenesis was quantified FIG. 8A as described above, and representative photomicrographs (6x) were taken with a stereomicroscope as shown in FIG. 8B. FIG. 8C shows a blot probed with an anti-Src antibody to confirm the expression of Src 251 in transfected cells as compared to mock treatments.

FIG. 9 illustrates the results of retroviral delivery of RCAS-Src 251 to human tumors. FIG. 9A is a micrograph that shows human medulloblastoma tumor fragment infected with RCAS-GFP (RCAS-Green Fluorescent Protein) expressing GFP exclusively in the tumor blood vessels (arrowhead) as detected by optical sectioning with a Bio Rad laser confocal scanning microscope (bar=500  $\mu$ m). FIG. 9B depicts data from tumors treated with topical application of retrovirus, which were allowed to grow for 3 or 6 days after which they were resected and wet weights determined. Data are expressed as the mean change in tumor weight (from the 50 mg tumor starting weight) +/- SEM of 2 replicates. FIG. 9C depicts in representative micrographs, medulloblastoma tumors surgically removed from the embryos (bar=350  $\mu$ m). The lower panels are high magnification views of each tumor showing the vasculature of each tumor in detail (bar=350  $\mu$ m). The arrowhead indicates blood vessel disruption in RCAS-Src251-treated tumors.

FIG. 10 is a diagram illustrating a restriction map of the RCASBP (RCAS) vector construct.

### Detailed Description of the Invention

#### A. Definitions

Amino Acid Residue: An amino acid formed upon chemical digestion (hydrolysis) of a polypeptide at its peptide linkages. The amino acid residues

described herein are preferably in the "L" isomeric form. However, residues in the "D" isomeric form can be substituted for any L-amino acid residue, as long as the desired functional property is retained by the polypeptide. NH<sub>2</sub> refers to the free amino group present at the amino terminus of a polypeptide. COOH refers to the free carboxy group present at the carboxy terminus of a polypeptide in keeping with standard polypeptide nomenclature (described in J. Biol. Chem., 243:3552-59 (1969) and adopted at 37 CFR §1.822(b)(2)).

It should be noted that all amino acid residue sequences are represented herein by formulae whose left and right orientation is in the conventional direction of amino-terminus to carboxy-terminus. Furthermore, a dash at the beginning or end of an amino acid residue sequence indicates a peptide bond to a further sequence of one or more amino acid residues.

Polypeptide: refers to a linear array of amino acid residues connected to one another by peptide bonds between the alpha-amino group and carboxy group of contiguous amino acid residues.

Peptide: as used herein refers to a linear array of no more than about 50 amino acid residues connected one to the other as in a polypeptide.

Cyclic peptide: refers to a compound having a ring structure that includes several amide bonds as in a typical peptide. The cyclic peptide can be a "head to tail" homodetic cyclic peptide, or it can contain a heterodetic ring structure in which the ring is closed by disulfide bridges, lactam bridges, thioesters, thioamides, guanidino, and the like linkages.

Protein: refers to a linear array of more than 50 amino acid residues connected one to the other as in a polypeptide.

Fusion protein: refers to a polypeptide containing at least two different polypeptide domains operatively linked by a typical peptide bond ("fused"), where the two domains correspond to peptides not found fused in nature.

Synthetic peptide: refers to a chemically produced chain of amino acid residues linked together by peptide bonds that is free of naturally occurring proteins and fragments thereof.

#### B. General Considerations

The present invention relates generally to the discovery that angiogenesis is mediated by the tyrosine kinase Src protein, and that angiogenesis can be



modulated by providing either active or inactive Src proteins for potentiating or inhibiting angiogenesis, respectively.

This discovery is important because of the role that angiogenesis, the formation of new blood vessels, plays in a variety of disease processes. Where tissues associated with a disease condition require angiogenesis for tissue growth, it is desirable to inhibit angiogenesis and thereby inhibit the diseased tissue growth. Where injured tissue requires angiogenesis for tissue growth and healing, it is desirable to potentiate or promote angiogenesis and thereby promote tissue healing and growth.

Where the growth of new blood vessels is the cause of, or contributes to, the pathology associated with a diseased tissue, inhibition of angiogenesis reduces the deleterious effects of the disease. By inhibiting angiogenesis, one can intervene in the disease, ameliorate the symptoms, and in some cases cure the disease.

Examples of tissue associated with disease and neovascularization that will benefit from inhibitory modulation of angiogenesis include rheumatoid arthritis, diabetic retinopathy, inflammatory diseases, restenosis, and the like. Where the growth of new blood vessels is required to support growth of a deleterious tissue, inhibition of angiogenesis will reduce the blood supply to the tissue and thereby contribute to reduction in tissue mass based on blood supply requirements.

Examples include growth of tumors where neovascularization is a continual requirement in order that the tumor grow beyond a few millimeters in thickness, and for the establishment of solid tumor metastases.

Where the growth of new blood vessels contributes to healing of tissue, potentiation of angiogenesis assists in healing. Examples include treatment of patients with ischemic limbs in which there is poor circulation in the limbs from diabetes or other conditions. Also contemplated for treatment are patients with chronic wounds that do not heal and therefore could benefit from the increase in vascular cell proliferation and neovascularization.

The methods of the present invention are effective in part because the therapy is highly selective for angiogenesis and not other biological processes.

As described earlier, angiogenesis includes a variety of processes involving neovascularization of a tissue including "sprouting", vasculogenesis, or vessel enlargement, all of which angiogenesis processes are effected by Src protein.

With the exception of traumatic wound healing, corpus luteum formation and embryogenesis, it is believed that the majority of angiogenesis processes are associated with disease processes. Accordingly, the present therapeutic methods are selective for the disease and do not have deleterious side effects.

5 C. Src Proteins

A tyrosine kinase Src protein for use in the present invention can vary depending upon the intended use. The terms "Src protein" or "Src" are used to refer to the various forms of tyrosine kinase Src proteins described herein, either in active or inactive forms.

10 An "active Src protein" refers to any of a variety of forms of src protein which potentiate angiogenesis. Assays to measure potentiation of angiogenesis are described herein, and are not to be construed as limiting. A protein is considered active if the level of angiogenesis is at least 10% greater, preferably 25% greater, and more preferably 50% greater than a control level where no src is added to the  
15 assay system. The preferred assay for measuring potentiation is the CAM assay using RCAS viral vector as described in the Examples in which the angiogenic index is calculated by counting branch points. A preferred active Src protein exhibits tyrosine kinase activity as well. Exemplary active Src proteins are described in the Examples, and include Src-A.

20 An "inactive Src protein" refers to any of a variety of forms of Src protein which inhibit angiogenesis. Assays to measure inhibition of angiogenesis are described herein, and are not to be construed as limiting. A protein is considered inactive if the level of angiogenesis is at least 10% lower, preferably 25% lower, and more preferably 50% lower than a control level where no exogenous Src is  
25 added to the assay system. The preferred assay for measuring inhibition is the CAM assay using RCAS viral vector as described in the Examples in which the angiogenic index is calculated by counting branch points. A preferred inactive Src protein exhibits reduced tyrosine kinase activity as well. Exemplary inactive Src proteins are described in the Examples, and include Src-251.

30 A Src protein useful in the present invention can be produced in any of a variety of methods including isolation from natural sources including tissue, production by recombinant DNA expression and purification, and the like. Src

protein can also be provided "in situ" by introduction of a gene therapy system to the tissue of interest which then expresses the protein in the tissue.

A gene encoding a Src protein can be prepared by a variety of methods known in the art, and the invention is not to be construed as limiting in this regard. For example, the natural history of Src is well known to include a variety of homologs from mammalian, avian, viral and the like species, and the gene can readily be cloned using cDNA cloning methods from any tissue expressing the protein. A preferred Src for use in the invention is a cellular protein, such as the mammalian or avian homologs designated c-Src. Particularly preferred is human c-Src.

10 D. Recombinant DNA Molecules and Expression Systems for Expression of a Src Protein

The invention describes several nucleotide sequences of particular use in the present invention. These sequences include sequences which encode a Src protein useful in the invention, and various DNA segments, recombinant DNA (rDNA) molecules and vectors constructed for expression of Src protein.

DNA molecules (segments) of this invention therefore can comprise sequences which encode whole structural genes, fragments of structural genes, and transcription units as described further herein.

A preferred DNA segment is a nucleotide sequence which encodes a Src protein as defined herein, or biologically active fragment thereof,

The amino acid residue sequence and nucleotide sequence of a preferred c-Src is described in the Examples.

A preferred DNA segment codes for an amino acid residue sequence substantially the same as, and preferably consisting essentially of, an amino acid residue sequence or portions thereof corresponding to a Src protein described herein. Representative and preferred DNA segments are further described in the Examples.

The amino acid residue sequence of a protein or polypeptide is directly related via the genetic code to the deoxyribonucleic acid (DNA) sequence of the structural gene that codes for the protein. Thus, a structural gene or DNA segment can be defined in terms of the amino acid residue sequence, i.e., protein or polypeptide, for which it codes.

An important and well known feature of the genetic code is its redundancy. That is, for most of the amino acids used to make proteins, more than one coding nucleotide triplet (codon) can code for or designate a particular amino acid residue. Therefore, a number of different nucleotide sequences may code for a particular amino acid residue sequence. Such nucleotide sequences are considered functionally equivalent since they can result in the production of the same amino acid residue sequence in all organisms. Occasionally, a methylated variant of a purine or pyrimidine may be incorporated into a given nucleotide sequence. However, such methylations do not affect the coding relationship in any way.

A nucleic acid is any polynucleotide or nucleic acid fragment, whether it be a polyribonucleotide or polydeoxyribonucleotide, i.e., RNA or DNA, or analogs thereof. In preferred embodiments, a nucleic acid molecule is in the form of a segment of duplex DNA, i.e., a DNA segment, although for certain molecular biological methodologies, single-stranded DNA or RNA is preferred.

DNA segments are produced by a number of means including chemical synthesis methods and recombinant approaches, preferably by cloning or by polymerase chain reaction (PCR). DNA segments that encode portions of a Src protein can easily be synthesized by chemical techniques, for example, the phosphotriester method of Matteucci et al, J. Am. Chem. Soc., 103:3185-3191, 1981, or using automated synthesis methods. In addition, larger DNA segments can readily be prepared by well known methods, such as synthesis of a group of oligonucleotides that define the DNA segment, followed by hybridization and ligation of oligonucleotides to build the complete segment. Alternative methods include isolation of a preferred DNA segment by PCR with a pair of oligonucleotide primers used on a cDNA library believed to contain members which encode a Src protein.

Of course, through chemical synthesis, any desired modifications can be made simply by substituting the appropriate bases for those encoding the native amino acid residue sequence. This method is well known, and can be readily applied to the production of the various different "modified" Src proteins described herein.

Furthermore, DNA segments consisting essentially of structural genes encoding a Src protein can be subsequently modified, as by site-directed or random mutagenesis, to introduce any desired substitutions.

### 1. Cloning a Src Gene

A Src gene can be cloned from a suitable source of genomic DNA or messenger RNA (mRNA) by a variety of biochemical methods. Cloning of these genes can be conducted according to the general methods described in the Examples and as known in the art.

Sources of nucleic acids for cloning a Src gene suitable for use in the methods of this invention can include genomic DNA or messenger RNA (mRNA) in the form of a cDNA library, from a tissue believed to express these proteins. A preferred tissue is human lung tissue, although any other suitable tissue may be used.

A preferred cloning method involves the preparation of a cDNA library using standard methods, and isolating the Src-encoding nucleotide sequence by PCR amplification using paired oligonucleotide primers based on the nucleotide sequences described herein. Alternatively, the desired cDNA clones can be identified and isolated from a cDNA or genomic library by conventional nucleic acid hybridization methods using a hybridization probe based on the nucleic acid sequences described herein. Other methods of isolating and cloning suitable src encoding nucleic acids are readily apparent to one skilled in the art.

### 2. Expression Vectors

A recombinant DNA molecule (rDNA) containing a DNA segment encoding a Src protein can be produced as described herein. In particular, an expressible rDNA can be produced by operatively (in frame, expressibly) linking a vector to a src encoding DNA segment. Thus, a recombinant DNA molecule is a hybrid DNA molecule comprising at least two nucleic acids of a nucleotide sequences not normally found together in nature.

The choice of vector to which the DNA segment is operatively linked depends directly, as is well known in the art, on the functional properties desired, e.g., protein expression, and the host cell to be transformed. A vector suitable for use in practicing the present invention is at least capable of directing the replication, and preferably also expression, of a structural gene included in the vector DNA segments to which it is operatively linked.

Both prokaryotic and eukaryotic expression vectors are familiar to one of ordinary skill in the art of vector construction, and are described by Ausubel, et

al., in Current Protocols in Molecular Biology, Wiley and Sons, New York (1993) and by Sambrook et al., Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory, (1989). These references also describe many of the general recombinant DNA methods referred to herein.

5 In one embodiment, a suitable vector includes a procaryotic replicon, i.e., a DNA sequence having the ability to direct autonomous replication and maintenance of the recombinant DNA molecule extrachromosomally in a procaryotic host cell, such as a bacterial host cell, transformed therewith. Such replicons are well known in the art. In addition, those embodiments that include a procaryotic  
10 replicon also include a gene whose expression confers drug resistance to a bacterial host transformed therewith. Typical bacterial drug resistance genes are those that confer resistance to ampicillin or tetracycline.

Those vectors that include a procaryotic replicon can also include a procaryotic promoter capable of directing the expression (transcription and  
15 translation) of a structural gene in a bacterial host cell, such as E. coli, transformed therewith. A promoter is an expression control element formed by a DNA sequence that permits binding of RNA polymerase and transcription to occur. Promoter sequences compatible with bacterial hosts are typically provided in plasmid vectors containing convenient restriction sites for insertion of a DNA  
20 segment of the present invention. Typical of such vector plasmids are pUC8, pUC9, pBR322 and pBR329 available from Biorad Laboratories, (Richmond, CA), pRSET available from Invitrogen (San Diego, CA) and pPL and pKK223 available from Pharmacia, Piscataway, N.J.

Expression vectors compatible with eukaryotic cells, preferably those  
25 compatible with vertebrate cells, can also be used to form the recombinant DNA molecules of the present invention. Eukaryotic cell expression vectors are well known in the art and are available from several commercial sources. Typically, such vectors are provided containing convenient restriction sites for insertion of the desired DNA segment. Typical of such vectors are pSVL and pKSV-10  
30 (Pharmacia), pBPV-1/pML2d (International Biotechnologies, Inc.), pTDT1 (ATCC, #31255), pRc/CMV (Invitrogen, Inc.), the preferred vector described in the Examples, and the like eukaryotic expression vectors.

A particularly preferred system for gene expression in the context of this invention includes a gene delivery component, that is, the ability to deliver the gene to the tissue of interest. Suitable vectors are "infectious" vectors such as recombinant DNA viruses, adenovirus or retrovirus vectors which are engineered to express the desired protein and have features which allow infection of preselected target tissues. Particularly preferred is the replication competent avian sarcoma virus (RCAS) described herein.

Mammalian cell systems that utilize recombinant viruses or viral elements to direct expression may be engineered. For example, when using adenovirus expression vectors, the coding sequence of a polypeptide may be ligated to an adenovirus transcription/ translation control complex, e.g., the late promoter and tripartite leader sequence. This chimeric gene may then be inserted into the adenovirus genome by in vitro or in vivo recombination. Insertion in a non-essential region of the viral genome (e.g., region E1 or E3) will result in a recombinant virus that is viable and capable of expressing the polypeptide in infected hosts (e.g., see Logan et al., Proc. Natl. Acad. Sci., USA, 81:3655-3659 (1984)). Alternatively, the vaccinia virus 7.5K promoter may be used (e.g., see, Mackett et al., Proc. Natl. Acad. Sci., USA, 79:7415-7419 (1982); Mackett et al., J. Virol., 49:857-864 (1984); Panicali et al., Proc. Natl. Acad. Sci., USA, 79:4927-4931 (1982)). Of particular interest are vectors based on bovine papilloma virus which have the ability to replicate as extrachromosomal elements (Sarver et al., Mol. Cell. Biol., 1:486 (1981)). Shortly after entry of this DNA into target cells, the plasmid replicates to about 100 to 200 copies per cell. Transcription of the inserted cDNA does not require integration of the plasmid into the host's chromosome, thereby yielding a high level of expression. These vectors can be used for stable expression by including a selectable marker in the plasmid, such as the neo gene. Alternatively, the retroviral genome can be modified for use as a vector capable of introducing and directing the expression of the polypeptide-encoding nucleotide sequence in host cells (Cone et al., Proc. Natl. Acad. Sci., USA, 81:6349-6353 (1984)). High level expression may also be achieved using inducible promoters, including, but not limited to, the metallothioneine IIA promoter and heat shock promoters.

Recently, long-term survival of cytomegalovirus (CMV) promoter versus Rous sarcoma virus (RSV) promoter-driven thymidine kinase (TK) gene therapy in nude mice bearing human ovarian cancer has been studied. Cell killing efficacy of adenovirus-mediated CMV promoter-driven herpes simplex virus TK gene therapy was found to be 2 to 10 time more effective than RSV driven therapy. (Tong et al., 1999, Hybridoma 18(1):93-97). The design of chimeric promoters for gene therapy applications, which call for low level expression followed by inducible high-level expression has also been described. (Suzuki et al., 1996, Human Gene Therapy 7:1883-1893).

For long-term, high-yield production of recombinant proteins, stable expression is preferred. Rather than using expression vectors which contain viral origins of replication, host cells can be transformed with a cDNA controlled by appropriate expression control elements (e.g., promoter and enhancer sequences, transcription terminators, polyadenylation sites, etc.), and a selectable marker. As mentioned above, the selectable marker in the recombinant plasmid confers resistance to the selection and allows cells to stably integrate the plasmid into their chromosomes and grow to form foci which in turn can be cloned and expanded into cell lines.

For example, following the introduction of foreign DNA, engineered cells may be allowed to grow for 1-2 days in an enriched media, and then are switched to a selective media. A number of selection systems may be used, including but not limited to the herpes simplex virus thymidine kinase (Wigler et al., Cell, 11:223 (1977)), hypoxanthine-guanine phosphoribosyltransferase (Szybalska et al, Proc. Natl. Acad. Sci., USA, 48:2026 (1962)), and adenine phosphoribosyltransferase (Lowy et al., Cell, 22:817 (1980)) genes, which can be employed in tk-, hgp<sup>r</sup>t- or ap<sup>r</sup>t- cells respectively. Also, antimetabolite resistance-conferring genes can be used as the basis of selection; for example, the genes for dhfr, which confers resistance to methotrexate (Wigler et al., Proc. Natl. Acad. Sci., USA, 77:3567 (1980); O'Hare et al., Proc. Natl. Acad. Sci., USA, 78:1527 (1981); gpt, which confers resistance to mycophenolic acid (Mulligan et al, Proc. Natl. Acad. Sci., USA, 78:2072, (1981)); neo, which confers resistance to the aminoglycoside G-418 (Colberre-Garapin et al, J. Mol. Biol., 150:1 (1981)); and hyg<sup>r</sup>, which confers resistance to hygromycin (Santerre et al, Gene,



30:147 (1984)). Recently, additional selectable genes have been described, namely trpB, which allows cells to utilize indole in place of tryptophan; hisD, which allows cells to utilize histinol in place of histidine (Hartman et al, Proc. Natl. Acad. Sci., USA, 85:804 (1988)); and ODC (ornithine decarboxylase) which  
5 confers resistance to the ornithine decarboxylase inhibitor, 2-(difluoromethyl)-DL-ornithine, DFMO (McConlogue L., In: Current Communications in Molecular Biology, Cold Spring Harbor Laboratory ed., (1987)).

The principal vectors contemplated for human gene therapy are derived from retroviral origin. (Wilson, 1997, Clin. Exp. Immunol. 107(Sup. 1):31-32; Bank et al., 1996, Bioessays 18(12):999-1007; Robbins et al., 1998, Pharmacol. Ther. 80(1):35-47). The therapeutic potential of gene transfer and antisense therapy has stimulated the development of many vector systems for treating a variety of tissues. (vasculature, Stephan et al., 1997, Fundam. Clin. Pharmacol. 11(2):97-  
15 110; Feldman et al., 1997, Cardiovasc. Res. 35(3):391-404; Vassalli et al., 1997, Cardiovasc. Res. 35(3):459-69; Baek et al., 1998, Circ. Res. 82(3):295-305; kidney, Lien et al., 1997, Kidney Int. Suppl. 61:S85-8; liver, Ferry et al., 1998, Hum Gene Ther. 9(14):1975-81; muscle, Marshall et al., 1998, Curr. Opin. Genet. Dev. 8(3):360-5). In addition to these tissues, a critical target for human gene  
20 therapy is cancer, either the tumor itself, or associated tissues. (Runnebaum, 1997, Anticancer Res. 17(4B):2887-90; Spear et al., 1998, J. Neurovirol. 4(2):133-47).

Specific examples of viral gene therapy vector systems readily adaptable for use in the methods of the present invention are briefly described below. Retroviral gene delivery has been recently reviewed by Federspiel and Hughes (1998, Methods in Cell Biol. 52:179-214) which describes in particular, the avian  
25 leukemia virus (ALV) retrovirus family (Federspiel et al., Proc. Natl. Acad. Sci., USA, 93:4931 (1996); Federspiel et al., Proc. Natl. Acad. Sci., USA, 91:11241 (1994)). Retroviral vectors, including ALV and murine leukemia virus (MLV) are further described by Svoboda (1998, Gene 206:153-163).

30 Modified retroviral/adenoviral expression systems can be readily adapted for practice of the methods of the present invention. For example, murine leukemia virus (MLV) systems are reviewed by Karavanas et al., 1998, Crit. Rev. in Oncology/Hematology 28:7-30. Adenovirus expression systems are reviewed by

Von Seggern and Nemerow in Gene Expression Systems (ed. Fernandez & Hoeffler, Academic Press, San Diego, CA, 1999, chapter 5, pages 112-157).

Protein expression systems have been demonstrated to have effective use both in vivo and in vitro. For example, efficient gene transfer to human squamous cell carcinomas by a herpes simplex virus (HSV) type 1 amplicon vector has been described. (Carew et al., 1998, Am. J. Surg. 176:404-408). Herpes simplex virus has been used for gene transfer to the nervous system. (Goins et al., 1997, J. Neurovirol. 3 (Sup. 1):S80-8). Targeted suicide vectors using HSV-TK has been tested on solid tumors. (Smiley et al., 1997, Hum. Gene Ther. 8(8):965-77). Herpes simplex virus type 1 vector has been used for cancer gene therapy on colon carcinoma cells. (Yoon et al., 1998, Ann. Surg. 228(3):366-74). Hybrid vectors have been developed to extend the length of time of transfection, including HSV/AAV (adeno-associated virus) hybrids for treating hepatocytes. (Fraefel et al., 1997, Mol. Med. 3(12):813-825).

Vaccinia virus has been developed for human gene therapy because of its large genome. (Peplinski et al., 1998, Surg. Oncol. Clin. N. Am. 7(3):575-88). Thymidine kinase-deleted vaccinia virus expressing purine nucleoside pyrophosphorylase has been described for use as a tumor directed gene therapy vector. (Puhlman et al., 1999, Human Gene Therapy 10:649-657).

Adeno-associated virus 2 (AAV) has been described for use in human gene therapy, however AAV requires a helper virus (such as adenovirus or herpes virus) for optimal replication and packaging in mammalian cells. (Snoeck et al., 1997, Exp. Nephrol. 5(6):514-20; Rabinowitz et al., 1998, Curr. Opin. Biotechnol. 9(5):470-5). However, in vitro packaging of an infectious recombinant AAV has been described, making this system much more promising. (Ding et al., 1997, Gene Therapy 4:1167-1172). It has been shown that the AAV mediated transfer of ecotropic retrovirus receptor cDNA allows ecotropic retroviral transduction of established and primary human cells. (Qing et al., 1997, J. Virology 71(7):5663-5667). Cancer gene therapy using an AAV vector expressing human wild-type p53 has been demonstrated. (Qazilbash et al., 1997, Gene Therapy 4:675-682). Gene transfer into vascular cells using AAV vectors has also been shown. (Maeda et al., 1997, Cardiovascular Res. 35:514-521). AAV has been demonstrated as a suitable vector for liver directed gene therapy. (Xiao et al., 1998, J. Virol. 72(12):10222-

6). AAV vectors have been demonstrated for use in gene therapy of brain tissues and the central nervous system. (Chamberlin et al., 1998, Brain Res. 793(1-2):169-75; During et al., 1998, Gene Therapy 5(6):820-7). AAV vectors have also been compared with adenovirus vectors (AdV) for gene therapy of the lung and transfer to human cystic fibrosis epithelial cells. (Teramoto et al., 1998, J. Virol. 72(11):8904-12).

Chimeric AdV/retroviral gene therapy vector systems which incorporate the useful qualities of each virus to create a nonintegrative AdV that is rendered functionally integrative via the intermediate generation of a retroviral producer cell. (Feng et al., 1997, Nat. Biotechnology 15(9):866-70; Bilbao et al., 1997, FASEB J 11(8):624-34). This powerful new generation of gene therapy vector has been adapted for targeted cancer gene therapy. (Bilbao et al., 1998, Adv. Exp. Med. Biol. 451:365-74). Single injection of AdV expressing p53 inhibited growth of subcutaneous tumor nodules of human prostate cancer cells. (Asgari et al., 1997, Int. J. Cancer 71(3):377-82). AdV mediated gene transfer of wild-type p53 in patients with advanced non-small cell lung cancer has been described. (Schuler et al., 1998, Human Gene Therapy 9:2075-2082). This same cancer has been the subject of p53 gene replacement therapy mediated by AdV vectors. (Roth et al., 1998, Semin. Oncol. 25(3 Suppl 8):33-7). AdV mediated gene transfer of p53 inhibits endothelial cell differentiation and angiogenesis in vivo. (Riccioni et al., 1998, Gene Ther. 5(6):747-54). Adenovirus-mediated expression of melanoma antigen gp75 as immunotherapy for metastatic melanoma has also been described. (Hirschowitz et al., 1998, Gene Therapy 5:975-983). AdV facilitates infection of human cells with ecotropic retrovirus and increases efficiency of retroviral infection. (Scott-Taylor, et al., 1998, Gene Ther. 5(5):621-9). AdV vectors have been used for gene transfer to vascular smooth muscle cells (Li et al., 1997, Chin. Med. J.(Engl) 110(12):950-4), squamous cell carcinoma cells (Goebel et al., 1998, Otolaryngol Head Neck Surg 119(4):331-6), esophageal cancer cells (Senmaru et al., 1998, Int J. Cancer 78(3):366-71), mesangial cells (Nahman et al., 1998, J. Investig. Med. 46(5):204-9), glial cells (Chen et al., 1998, Cancer Res. 58(16):3504-7), and to the joints of animals (Ikeda et al., 1998, J. Rheumatol. 25(9):1666-73). More recently, catheter-based pericardial gene transfer mediated by AcV vectors has been demonstrated. (March et al., 1999, Clin. Cardiol. 22(1

Suppl 1):123-9). Manipulation of the AdV system with the proper controlling genetic elements allows for the AdV-mediated regulable target gene expression in vivo. (Burcin et al., 1999, PNAS (USA) 96(2):355-60).

5 Alphavirus vectors have been developed for human gene therapy applications, with packaging cell lines suitable for transformation with expression cassettes suitable for use with Sindbis virus and Semliki Forest virus-derived vectors. (Polo et al., 1999, Proc. Natl. Acad. Sci., USA, 96:4598-4603). Noncytopathic flavivirus replicon RNA-based systems have also been developed. (Varnavski et al., 1999, Virology 255(2):366-75). Suicide HSV-TK gene containing sinbis virus  
10 vectors have been used for cell-specific targeting into tumor cells. (Iijima et al., 1998, Int. J. Cancer 80(1):110-8).

Retroviral vectors based on human foamy virus (HFV) also show promise as gene therapy vectors. (Trobridge et al., 1998, Human Gene Therapy 9:2517-2525). Foamy virus vectors have been designed for suicide gene therapy. (Nestler et al., 1997, Gene Ther. 4(11):1270-7). Recombinant murine cytomegalovirus and  
15 promoter systems have also been used as vectors for high level expression. (Manning et al., 1998, J. Virol. Meth. 73(1):31-9; Tong et al., 1998, Hybridoma 18(1):93-7).

Gene delivery into non-dividing cells has been made feasible by the generation  
20 of Sendai virus based vectors. (Nakanishi et al., 1998, J. Controlled Release 54(1):61-8).

In other efforts to enable the transformation of non-dividing somatic cells, lentiviral vectors have been explored. Gene therapy of cystic fibrosis using a replication-defective human immunodeficiency virus (HIV) based vector has been  
25 described. (Goldman et al., 1997, Human Gene Therapy 8:2261-2268). Sustained expression of genes delivered into liver and muscle by lentiviral vectors has also been shown. (Kafri et al., 1997, Nat. Genet. 17(3):314-7). However, safety concerns are predominant, and improved vector development is proceeding rapidly. (Kim et al., 1998, J. Virol. 72(2):994-1004). Examination of the HIV  
30 LTR and Tat yield important information about the organization of the genome for developing vectors. (Sadaie et al., 1998, J. Med. Virol. 54(2):118-28). Thus the genetic requirements for an effective HIV based vector are now better understood. (Gasmi et al., 1999, J. Virol. 73(3):1828-34). Self inactivating vectors, or

conditional packaging cell lines have been described. (for example Zuffery et al., 1998, J. Virol. 72(12):9873-80; Miyoshi et al., 1998, J. Virol. 72(10):8150-7; Dull et al., 1998, J. Virol. 72(11):8463-71; and Kaul et al., 1998, Virology 249(1):167-74). Efficient transduction of human lymphocytes and CD34+ cells by HIV vectors has been shown. (Douglas et al., 1999, Hum. Gene Ther. 10(6):935-45; Miyoshi et al., 1999, Science 283(5402):682-6). Efficient transduction of nondividing human cells by feline immunodeficiency virus (FIV) lentiviral vectors has been described, which minimizes safety concerns with using HIV based vectors. (Poeschla et al., 1998, Nature Medicine 4(3):354-357). Productive infection of human blood mononuclear cells by FIV vectors has been shown. (Johnston et al., 1999, J. Virol. 73(3):2491-8).

While many viral vectors are difficult to handle, and capacity for inserted DNA limited, these limitations and disadvantages have been addressed. For example, in addition to simplified viral packaging cell lines, Mini-viral vectors, derived from human herpes virus, herpes simplex virus type 1 (HSV-1), and Epstein-Barr virus (EBV), have been developed to simplify manipulation of genetic material and generation of viral vectors. (Wang et al., 1996, J. Virology 70(12):8422-8430). Adaptor plasmids have been previously shown to simplify insertion of foreign DNA into helper-independent Retroviral vectors. (1987, J. Virology 61(10):3004-3012).

Viral vectors are not the only means for effecting gene therapy, as several non-viral vectors have also been described. A targeted non-viral gene delivery vector based on the use of Epidermal Growth Factor/DNA polyplex (EGF/DNA) has been shown to result in efficient and specific gene delivery. (Cristiano, 1998, Anticancer Res. 18:3241-3246). Gene therapy of the vasculature and CNS have been demonstrated using cationic liposomes. (Yang et al., 1997, J. Neurotrauma 14(5):281-97). Transient gene therapy of pancreatitis has also been accomplished using cationic liposomes. (Denham et al., 1998, Ann. Surg. 227(6):812-20). A chitosan-based vector/DNA complexes for gene delivery have been shown to be effective. (Erbacher et al., 1998, Pharm. Res. 15(9):1332-9). A non-viral DNA delivery vector based on a terplex system has been described. (Kim et al., 1998, 53(1-3):175-82). Virus particle coated liposome complexes have also been used to

effect gene transfer. (Hirai et al., 1997, Biochem. Biophys. Res. Commun. 241(1):112-8).

5 Cancer gene therapy by direct tumor injections of nonviral T7 vector encoding a thymidine kinase gene has been demonstrated. (Chen et al., 1998, Human Gene Therapy 9:729-736). Plasmid DNA preparation is important for direct injection gene transfer. (Horn et al., 1995, Hum. Gene Ther. 6(5):656-73). Modified plasmid vectors have been adapted specifically for direct injection. (Hartikka et al., 1996, Hum. Gene Ther. 7(10):1205-17).

10 Thus, a wide variety of gene transfer/gene therapy vectors and constructs are known in the art. These vectors are readily adapted for use in the methods of the present invention. By the appropriate manipulation using recombinant DNA/molecular biology techniques to insert an operatively linked src (either active or inactive) into the selected expression/delivery vector, many equivalent vectors for the practice of the present invention can be generated.

15 E. Methods For Modulation of Angiogenesis

The invention provides for a method for the modulation of angiogenesis in a tissue associated with a disease process or condition, and thereby effect events in the tissue which depend upon angiogenesis. Generally, the method comprises administering to the tissue associated with a disease process or condition, a composition comprising an angiogenesis-modulating amount of a Src protein or nucleic acid vector expressing active or inactive Src.

20 As described herein, any of a variety of tissues, or organs comprised of organized tissues, can support angiogenesis in disease conditions including skin, muscle, gut, connective tissue, joints, bones and the like tissue in which blood vessels can invade upon angiogenic stimuli.

25 The patient treated according to the present invention in its many embodiments is desirably a human patient, although it is to be understood that the principles of the invention indicate that the invention is effective with respect to all mammals, which are intended to be included in the term "patient". In this context, a mammal is understood to include any mammalian species in which treatment of tissue associated with diseases involving angiogenesis is desirable, particularly agricultural and domestic mammalian species.

Thus the method comprises administering to a patient a therapeutically effective amount of a physiologically tolerable composition containing a Src protein or DNA vector for expressing a Src protein in practicing the methods of the invention.

5       The dosage ranges for the administration of a Src protein depend upon the form of the protein, and its potency, as described further herein. The dosage amounts are large enough to produce the desired effect in which angiogenesis and the disease symptoms mediated by angiogenesis are ameliorated. The dosage should not be so large as to cause adverse side effects, however, such as  
10   hyperviscosity syndromes, pulmonary edema, congestive heart failure, and the like. Generally, the dosage will vary with the age, condition, sex of the patient, and extent of the disease in the patient, and can be readily determined by one of skill in the art. The dosage can also be adjusted by the individual physician in the event of any complication.

15       A therapeutically effective amount is an amount of Src protein, or nucleic acid encoding for (active or inactive) src protein, sufficient to produce a detectable modulation of angiogenesis in the tissue being treated, ie., an angiogenesis-modulating amount. Modulation of angiogenesis can be measured by CAM assay as described herein, or by other methods known to one skilled in the art.

20       The Src protein or nucleic acid vector expressing the Src protein can be administered parenterally by injection or by gradual infusion over time. Although the tissue to be treated can typically be accessed in the body by systemic administration, and therefore is most often treated by intravenous administration of therapeutic compositions, other tissues and delivery means are contemplated as  
25   well. Thus, compositions of the invention can be administered intravenously, intraperitoneally, intramuscularly, subcutaneously, intracavity, transdermally, and can be delivered by peristaltic means.

      The therapeutic compositions containing a Src protein or nucleic acid vector expressing the Src protein can be conventionally administered intravenously, as by  
30   injection of a unit dose, for example. The term "unit dose" when used in reference to a therapeutic composition of the present invention refers to physically discrete units suitable as unitary dosage for the subject, each unit containing a

predetermined quantity of active material calculated to produce the desired therapeutic effect in association with the required diluent; i.e., carrier, or vehicle.

In one preferred embodiment the reagent is administered in a single dosage intravenously. Localized administration can be accomplished by direct injection or  
5 by taking advantage of anatomically isolated compartments, isolating the microcirculation of target organ systems, reperfusion in a circulating system, or catheter based temporary occlusion of target regions of vasculature associated with diseased tissues.

The compositions are administered in a manner compatible with the dosage  
10 formulation, and in a therapeutically effective amount. The quantity to be administered and timing depends on the subject to be treated, capacity of the subject's system to utilize the active ingredient, and degree of therapeutic effect desired. Precise amounts of active ingredient required to be administered depend  
15 on the judgement of the practitioner and are peculiar to each individual. However, suitable dosage ranges for systemic application are disclosed herein and depend on the route of administration. Suitable regimes for administration are also variable, but are typified by an initial administration followed by repeated doses at one or more hour intervals by a subsequent injection or other administration.

Alternatively, continuous intravenous infusion sufficient to maintain concentrations  
20 in the blood in the ranges specified for in vivo therapies are contemplated.

1. Inhibition of Angiogenesis

Inhibition of angiogenesis is important in a variety of diseases, referred to as angiogenic diseases. Such diseases include, but are not limited to, inflammatory disorders such as immune and non-immune inflammation, chronic articular  
25 rheumatism and psoriasis, disorders associated with inappropriate or inopportune invasion of vessels such as diabetic retinopathy, neovascular glaucoma, restenosis, capillary proliferation in atherosclerotic plaques and osteoporosis, and cancer associated disorders, such as solid tumors, solid tumor metastases, angiofibromas, retrolental fibroplasia, hemangiomas, Kaposi sarcoma and the like cancers which  
30 require neovascularization to support tumor growth.

Thus, methods which inhibit angiogenesis in a tissue associated with a disease condition ameliorates symptoms of the disease and, depending upon the disease, can contribute to cure of the disease. In one embodiment, the invention



contemplates inhibition of angiogenesis, per se, in a tissue associated with a disease condition. The extent of angiogenesis in a tissue, and therefore the extent of inhibition achieved by the present methods, can be evaluated by a variety of methods.

5        Thus, in one related embodiment, a tissue to be treated is an inflamed tissue and the angiogenesis to be inhibited is inflamed tissue angiogenesis where there is neovascularization of inflamed tissue. In this class the method contemplates inhibition of angiogenesis in arthritic tissues, such as in a patient with chronic articular rheumatism, in immune or non-immune inflamed tissues, in psoriatic  
10       tissue and the like.

         In another related embodiment, a tissue to be treated is a retinal tissue of a patient with a retinal disease such as diabetic retinopathy, macular degeneration or neovascular glaucoma and the angiogenesis to be inhibited is retinal tissue angiogenesis where there is neovascularization of retinal tissue.

15       In an additional related embodiment, a tissue to be treated is a tumor tissue of a patient with a solid tumor, a metastases, a skin cancer, a breast cancer, a hemangioma or angiofibroma and the like cancer, and the angiogenesis to be inhibited is tumor tissue angiogenesis where there is neovascularization of a tumor tissue. Typical solid tumor tissues treatable by the present methods include lung,  
20       pancreas, breast, colon, laryngeal, ovarian, and the like tissues. Inhibition of tumor tissue angiogenesis is a particularly preferred embodiment because of the important role neovascularization plays in tumor growth. In the absence of neovascularization of tumor tissue, the tumor tissue does not obtain the required nutrients, slows in growth, ceases additional growth, regresses and ultimately  
25       becomes necrotic resulting in killing of the tumor.

         Stated in other words, the present invention provides for a method of inhibiting tumor neovascularization by inhibiting tumor angiogenesis according to the present methods. Similarly, the invention provides a method of inhibiting tumor growth by practicing the angiogenesis-inhibiting methods.

30       The methods are also particularly effective against the formation of metastases because (1) their formation requires vascularization of a primary tumor so that the metastatic cancer cells can exit the primary tumor and (2) their establishment in a secondary site requires neovascularization to support growth of the metastases.

In a related embodiment, the invention contemplates the practice of the method in conjunction with other therapies such as conventional chemotherapy directed against solid tumors and for control of establishment of metastases. The administration of angiogenesis inhibitor is typically conducted during or after  
5 chemotherapy, although it is preferably to inhibit angiogenesis after a regimen of chemotherapy at times where the tumor tissue will be responding to the toxic assault by inducing angiogenesis to recover by the provision of a blood supply and nutrients to the tumor tissue. In addition, it is preferred to administer the angiogenesis inhibition methods after surgery where solid tumors have been  
10 removed as a prophylaxis against metastases.

Insofar as the present methods apply to inhibition of tumor neovascularization, the methods can also apply to inhibition of tumor tissue growth, to inhibition of tumor metastases formation, and to regression of established tumors.

Restenosis is a process of smooth muscle cell (SMC) migration and  
15 proliferation into the tissue at the site of percutaneous transluminal coronary angioplasty which hampers the success of angioplasty. The migration and proliferation of SMC's during restenosis can be considered a process of angiogenesis which is inhibited by the present methods. Therefore, the invention also contemplates inhibition of restenosis by inhibiting angiogenesis according to  
20 the present methods in a patient following angioplasty procedures. For inhibition of restenosis, the inactivated tyrosine kinase is typically administered after the angioplasty procedure because the coronary vessel wall is at risk of restenosis, typically for from about 2 to about 28 days, and more typically for about the first 14 days following the procedure.

25 The present method for inhibiting angiogenesis in a tissue associated with a disease condition, and therefore for also practicing the methods for treatment of angiogenesis-related diseases, comprises contacting a tissue in which angiogenesis is occurring, or is at risk for occurring, with a composition comprising a therapeutically effective amount of an inactivated Src protein or vector expressing the protein.  
30

Inhibition of angiogenesis and tumor regression occurs as early as 7 days after the initial contacting with the therapeutic composition. Additional or prolonged

exposure to inactive Src protein is preferable for 7 days to 6 weeks, preferably about 14 to 28 days.

## 2. Potentialiation of Angiogenesis

In cases where it is desirable to promote or potentiate angiogenesis, administration of an active Src protein to the tissue is useful. The routes and timing of administration are comparable to the methods described hereinabove for inhibition.

## F. Therapeutic Compositions

The present invention contemplates therapeutic compositions useful for practicing the therapeutic methods described herein. Therapeutic compositions of the present invention contain a physiologically tolerable carrier together with a Src protein or vector capable of expressing a Src protein as described herein, dissolved or dispersed therein as an active ingredient. In a preferred embodiment, the therapeutic composition is not immunogenic when administered to a mammal or human patient for therapeutic purposes.

As used herein, the terms "pharmaceutically acceptable", "physiologically tolerable" and grammatical variations thereof, as they refer to compositions, carriers, diluents and reagents, are used interchangeably and represent that the materials are capable of administration to or upon a mammal without the production of undesirable physiological effects such as nausea, dizziness, gastric upset and the like.

The preparation of a pharmacological composition that contains active ingredients dissolved or dispersed therein is well understood in the art and need not be limited based on formulation. Typically such compositions are prepared as injectable either as liquid solutions or suspensions, however, solid forms suitable for solution, or suspensions, in liquid prior to use can also be prepared. The preparation can also be emulsified or presented as a liposome composition.

The active ingredient can be mixed with excipients which are pharmaceutically acceptable and compatible with the active ingredient and in amounts suitable for use in the therapeutic methods described herein. Suitable excipients are, for example, water, saline, dextrose, glycerol, ethanol or the like and combinations thereof. In addition, if desired, the composition can contain minor amounts of

auxiliary substances such as wetting or emulsifying agents, pH buffering agents and the like which enhance the effectiveness of the active ingredient.

The therapeutic composition of the present invention can include pharmaceutically acceptable salts of any salt-forming components therein.

5 Pharmaceutically acceptable salts include the acid addition salts (formed with the free amino groups of the polypeptide) that are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids as acetic, tartaric, mandelic and the like. Salts formed with the free carboxyl groups can also be derived from inorganic bases such as, for example, sodium, potassium,  
10 ammonium, calcium or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, 2-ethylamino ethanol, histidine, procaine and the like.

Physiologically tolerable carriers for the active ingredients are well known in the art. Exemplary of liquid carriers are sterile aqueous solutions that contain no  
15 materials in addition to the active ingredients and water, or contain a buffer such as sodium phosphate at physiological pH value, physiological saline or both, such as phosphate-buffered saline. Still further, aqueous carriers can contain more than one buffer salt, as well as salts such as sodium and potassium chlorides, dextrose, polyethylene glycol and other solutes.

20 Liquid compositions can also contain liquid phases in addition to and to the exclusion of water. Exemplary of such additional liquid phases are glycerin, vegetable oils such as cottonseed oil, and water-oil emulsions.

A therapeutic composition contains an angiogenesis-modulating amount of an Src protein of the present invention, or sufficient recombinant DNA expression  
25 vector to express an effective amount of Src protein, typically formulated to contain an amount of at least 0.1 weight percent of Src protein per weight of total therapeutic composition. A weight percent is a ratio by weight of Src protein to total composition. Thus, for example, 0.1 weight percent is 0.1 grams of Src protein per 100 grams of total composition. For DNA expression vectors, the  
30 amount administered depends on the properties of the expression vector, the tissue to be treated, and the like considerations.

### G. Article of Manufacture

The invention also contemplates an article of manufacture which is a labelled container for providing a Src protein of the invention. An article of manufacture comprises packaging material provided with appropriate labeling for the disease  
5 condition to be treated and a pharmaceutical agent contained within the packaging material.

The pharmaceutical agent in an article of manufacture is any of the compositions of the present invention suitable for providing a Src protein and formulated into a pharmaceutically acceptable form as described herein according  
10 to the disclosed indications. Thus, the composition can comprise a Src protein or a DNA molecule which is capable of expressing a Src protein. The article of manufacture contains an amount of pharmaceutical agent sufficient for use in treating a condition indicated herein, either in unit or multiple dosages.

The packaging material comprises a label which indicates the use of the  
15 pharmaceutical agent contained therein, e.g., for treating conditions assisted by the inhibition or potentiation of angiogenesis, and the like conditions disclosed herein. The label can further include instructions for use and related information as may be required for marketing. The packaging material can include container(s) for storage of the pharmaceutical agent.

As used herein, the term packaging material refers to a material such as glass,  
20 plastic, paper, foil, and the like capable of holding within fixed means a pharmaceutical agent. Thus, for example, the packaging material can be plastic or glass vials, laminated envelopes and the like containers used to contain a pharmaceutical composition including the pharmaceutical agent.

25 In preferred embodiments, the packaging material includes a label that is a tangible expression describing the contents of the article of manufacture and the use of the pharmaceutical agent contained therein.

### Examples

30 The following examples relating to this invention are illustrative and should not, of course, be construed as specifically limiting the invention. Moreover, such variations of the invention, now known or later developed, which would be within

the purview of one skilled in the art are to be considered to fall within the scope of the present invention hereinafter claimed.

1. Preparation of c-Src Expression Constructs

5 For preparing the expression constructs useful in modulating angiogenesis by the methods of the present invention, c-Src cDNA is manipulated and inserted into an expression construct/vector.

10 The cDNA sequence encoding for wild-type (i.e., endogenous) chicken c-Src is shown in FIG. 1 (SEQ ID NO.:2) with the encoded amino acid residue sequence shown in FIG. 2 (SEQ ID NO.:3). The encoded protein sequence is translated from the cDNA nucleotide positions 112 to 1713. The nucleic acid sequence corresponding to the nucleic acid sequence of human c-Src cDNA (SEQ ID NO.:4) and encoded amino acid residue (SEQ ID NO.:5) sequences are shown respectively in FIGs. 3 and 4. For the human protein sequence, the coding  
15 sequence begins at nucleotide position 134 to 1486 of the cDNA.

Wild-type as well as a number of mutated c-Src cDNAs were prepared. Mutated c-Src constructs were prepared by site-directed mutagenesis as described by Kaplan et al., EMBO J., 13:4745-4756 (1994). The mutated c-Src constructs for encoding mutated c-Src proteins for use in the methods of the present invention  
20 are described in Kaplan et al., id.. Kaplan et al. describe various mutated c-Src constructs and encoded proteins of useful for the practice of this invention. For example, Kaplan et al. depict several products of chicken c-src alleles in their FIG. 1, including SrcA and Src251.

Two categories of c-Src function to modulate angiogenesis are described. As  
25 previously discussed, one category contains Src molecules that increase angiogenesis and thus are considered to be active proteins. Wild-type Src along with various mutations are shown in the present invention to induce angiogenesis. One preferred mutation of wild type c-src which functions in this context with respect to its ability to induce blood vessel growth and therefore increase tumor  
30 weight in vivo is the Src A mutant having a point mutation at amino acid (aa) residue position 527 changing tyrosine 527 to phenylalanine. This site is normally a site for negative regulation by the c-Src kinase, referred to as kinase CSK. When CSK phosphorylates aa527 in the wild-type src, the protein is inactivated.

However, in mutated Src A, the regulatory tyrosine converted to phenylalanine thus conferring upon the protein a constitutively (i.e., permanently) active protein not subject to inactivation by phosphorylation.

5 Mutations in src have also been shown to have the opposite modulatory effect on angiogenesis, inhibiting angiogenesis instead of stimulating it. Such mutations are referred to as inactive src mutations. Proteins having mutation that confer this inhibitory activity are also referred to as dominant negative Src proteins in that they inhibit neovascularization, including that resulting from endogenous activity of Src as well as enhanced Src activity resulting from growth factor stimulation.

10 Thus certain mutations of wild type c-src of the present invention can also function as a dominant negative with respect to their ability to block blood vessel growth, and for example, therefore decrease tumor weight in vivo.

Such preferred inhibitory c-Src protein includes the Src 251 in which only the first 251 amino acids of Src are expressed. This construct lacks the entire kinase

15 domain and is therefore referred to as "kinase dead" src protein. A second construct is the Src (K295M) mutation in which the lysine amino acid residue 295 is mutated into a methionine. This point mutation in the kinase domain prevents ATP binding and also blocks kinase-dependent Src functions related to vascular cell and tumor cell signaling and proliferation.

20 For example, for the mutation at residue 527, as long as the resultant mutated amino acid residue is not tyrosine, serine, or threonine, the present invention contemplates that the presence of an alternate amino acid at the desired position will result in a Src protein with a desired active, angiogenesis promoting modulatory activity.

25 With respect to the point mutations, any mutation resulting in the desired inhibitory or stimulatory activity is contemplated for use in this invention. Fusion protein constructs combining the desired src protein (mutation or fragment thereof) with expressed amino acid tags, antigenic epitopes, fluorescent protein, or other such protein or peptides are also contemplated, so long as the desired modulating

30 effect of the src protein is intact.

**TABLE I**

<u>Src/Mutation</u>	<u>Src Function</u>	<u>Effect on Angiogenesis</u>
c-Src	+	active stimulates
SrcA (T527F)	+	active stimulates
5 Src527(point)	+	active stimulates
Src251	-	inactive inhibits
Src (truncate)	-	inactive inhibits
Src(K295M)	-	inactive inhibits
Src295 (point)	-	inactive inhibits

10

One preferred expression construct for use in the present invention is the RCASBP(A) construct (SEQ ID NO.:1). This expression vector is based on a series of replication competent avian sarcoma viruses with an enhanced Bryan polymerase (BP) for improved titre, and is specific for the A type envelope glycoprotein expressed on normal avian cells (Reviewed in Methods in Cell Biology, 52:179-214 (1997); see also, Hughes et al., 1987, J. Virol. 61:3004-3012; Fekete & Cepko, 1993, Mol. Cellular Biol. 13(4):2604-2613; Itoh et al., 1996, Development 122:291-300; and Stott et al., 1998, BioTechniques 24:660-666). The complete sequence of RCASBP(A) (SEQ ID NO.:1) is given in the attached sequence listing, and a restriction map of the construct is depicted as FIG. 10, referred to herein as RCAS.

20

The original Src 251 construct was subcloned by Dr. Pam Schwartzberg, at NIH in Dr. Harold Varmus' laboratory. Briefly, cloning of a src cDNA sequence for expression thereof was accomplished by inserting a linker containing Not I-BstB1-Not I restriction sites into a unique Not I site in the 5' end of Src 251. Src has a unique Cla I site at the 3' end. Digestion of Src 251 with BstB1 and Cla I generated a BstB1-ClaI fragment which was then ligated into the Cla I site on RCASBP(A). A BstB1 overhang allows for ligation with a Cla I overhang that will not be recut with Cla I. The src constructs suitable for use in practicing the present invention are readily obtained in the above vector by first digesting the RCAS vector containing Src 251 with Not I and Cla I (in a DAM+ background) to allow for insertion of a similarly digested Src cDNA. Therefore this initial RCASBP(A) construct containing Src 251 was further used to subclone all other

25

30



Src constructs as described above and in Kaplan et al. (1994, The EMBO J. 13(20):4745-4756), into RCASBP(A) via a Not I-Cla I fragment generated through the Src 251 construction. To produce the desired c-src mutations in the cDNA, standard site-directed mutagenesis procedures familiar to one of ordinary skill in the art were utilized. PCR primers designed to incorporate the desired mutations were also designed with restriction sites to facilitate subsequent cloning steps. Entire segments of Src encoding nucleic acid sequences are deleted from the nucleic acid constructs through PCR amplification techniques based on the known cDNA sequences of chicken, human and the like homologs of Src and subsequent formation of new constructs.

In one embodiment of the invention, the 3' PCR primer used to amplify src nucleic acids also encodes for an in-frame sequence. Use of this primer adds a 9E10-myc epitope tag to the carboxyl terminus of the subsequent Src construct.

The following amino acids were added after amino acid 251 of Src to generate vector constructs containing the 9E10-myc epitope tag: VDMEQKLIAEEDLN (SEQ ID NO.: 6). Two separate PCRs were carried out for each construct and similar results were obtained. All mutant constructs constructed by PCR were also sequenced by PCR to confirm predicted DNA sequence of clones. Wild-type and mutated Src cDNAs for use in the expression systems of the present invention are also available from Upstate Biotech Laboratories, Lake Placid, NY which sells avian as well as human src, and several kinase dead and activated mutated forms.

Alternative expression vectors for use in the expressing the Src proteins of the present invention also include adenoviral vectors as described in US Patent Numbers 4,797,368, 5,173,414, 5,436,146, 5,589,377, and 5,670,488.

Alternative methods for the delivery of the Src modulatory proteins include delivery of the Src cDNA with a non-viral vector system as described in US Patent Number 5,675,954 and delivery of the cDNA itself as naked DNA as described in US Patent Number 5,589,466. Delivery of constructs of this invention is also not limited to topical application of a viral vector as described in the CAM assay system below. For example, viral vector preparations are also injected intravenously for systemic delivery into the vascular bed. These vectors are also targetable to sites of increased neovascularization by localized injection of a tumor, as an example.

In vitro expressed proteins are also contemplated for delivery thereof following expression and purification of the selected Src protein by methods useful for delivery of proteins or polypeptides. One such method includes liposome delivery systems, such as described in US Patent Numbers 4,356,167, 5,580,575, 5,542,935 and 5,643,599. Other vector and protein delivery systems are well known to those of ordinary skill in the art for use in the expression and/or delivery of the Src proteins of the present invention.

2. Characterization of the Untreated Chick Chorioallantoic Membrane (CAM)

A. Preparation of the CAM

Angiogenesis can be induced on the chick chorioallantoic membrane (CAM) after normal embryonic angiogenesis has resulted in the formation of mature blood vessels. Angiogenesis has been shown to be induced in response to specific cytokines or tumor fragments as described by Leibovich et al., Nature, 329:630 (1987) and Ausprunk et al., Am. J. Pathol., 79:597 (1975). CAMs were prepared from chick embryos for subsequent induction of angiogenesis and inhibition thereof. Ten day old chick embryos were obtained from McIntyre Poultry (Lakeside, CA) and incubated at 37°C with 60% humidity. A small hole was made through the shell at the end of the egg directly over the air sac with the use of a small crafts drill (Dremel, Division of Emerson Electric Co. Racine WI). A second hole was drilled on the broad side of the egg in a region devoid of embryonic blood vessels determined previously by candling the egg. Negative pressure was applied to the original hole, which resulted in the CAM (chorioallantoic membrane) pulling away from the shell membrane and creating a false air sac over the CAM. A 1.0 centimeter (cm) x 1.0 cm square window was cut through the shell over the dropped CAM with the use of a small model grinding wheel (Dremel). The small window allowed direct access to the underlying CAM.

The resultant CAM preparation was then either used at 6 days of embryogenesis, a stage marked by active neovascularization, without additional treatment to the CAM reflecting the model used for evaluating effects on embryonic neovascularization or used at 10 days of embryogenesis where angiogenesis has subsided. The latter preparation was thus used in this invention

for inducing renewed angiogenesis in response to cytokine treatment or tumor contact as described below.

### 3. CAM Angiogenesis Assay

#### A. Angiogenesis Induced by Growth Factors

5 Angiogenesis has been shown to be induced by cytokines or growth factors.

Angiogenesis was induced by placing a 5 millimeter (mm) X 5 mm Whatman filter disk (Whatman Filter paper No.1) saturated with Hanks Balanced Salt Solution (HBSS, GIBCO, Grand Island, NY) or HBSS containing 2 micrograms/milliliter ( $\mu\text{g/ml}$ ) recombinant basic fibroblast growth factor (bFGF) or vascular endothelial cell growth factor (VEGF) (Genzyme, Cambridge, MA) on the CAM of either a 9 or 10 day chick embryo in a region devoid of blood vessels and the windows were latter sealed with tape. Other concentrations of growth factors are also effective at inducing blood vessel growth. For assays where inhibition of angiogenesis is evaluated with intravenous injections of antagonists, angiogenesis is first induced with 1-2  $\mu\text{g/ml}$  bFGF or VEGF in fibroblast growth medium. Angiogenesis was monitored by photomicroscopy after 72 hours.

#### B. Embryonic Angiogenesis

The CAM preparation for evaluating the effect of angiogenesis inhibitors on the natural formation of embryonic neovasculature is the 6 day embryonic chick embryo as previously described. At this stage in development, the blood vessels are undergoing de novo growth and thus provides a useful system for assessing angiogenesis modulation by the Src proteins of the present invention. The CAM system is prepared as described above with the exception that the assay is performed at embryonic day 6 rather than at day 9 or 10.

### 25 4. Modulation of Angiogenesis as Measured in the CAM Assay

To assess the effect of Src proteins on angiogenesis, the following assays were performed on 10 day old chick CAM preparations. Five  $\mu\text{g}$  of RCAS constructs prepared as described in Example 1 were transfected into the chicken immortalized fibroblast line, DF-1 (gift of Doug Foster, U. of Minn.). This cell line as well as primary chick embryo fibroblasts were capable of producing virus, however the DF-1 cell line produced higher titres. Viral supernatants were collected from subconfluent DF-1 producer cell lines in serum free CLM media [composition: F-10 media base supplemented with DMSO, folic acid, glutamic acid, and MEM

vitamin solution]. Thirty-five ml of viral supernatant were concentrated by ultracentrifugation at 4°C for 2 hours at 22,000 rpm. These concentrated viral pellets were resuspended in 1/100 the original volume in serum-free CLM media, aliquoted and stored at -80°C. The titre was assessed by serial dilution of a control viral vector having a nucleotide sequence encoding green fluorescent protein (GFP), referred to as RCAS-GFP, infection on primary chick embryo fibroblasts that were incubated for 48-72 hours. The titres of viral stock that were obtained following concentration routinely exceeded 10<sup>8</sup> I.u./ml. For the CAM assay using the viral stocks, cortisone acetate soaked Whatman filter disks 6 mm in diameter were prepared in 3 mg/ml cortisone acetate for 30 minutes in 95% ethanol. The disks were dried in a laminar flow hood and then soaked on 20 µl of viral stock per disk for 10 minutes. These disks were applied to the CAM of 9 or 10 day chick embryos and sealed with cellophane tape and incubated at 37°C for 18-24 hr. Then either mock PBS or growth factors were added at a concentration of 5 µg/ml to the CAM in a 20 µl volume of the appropriate virus stock as an additional boost of virus to the CAM tissue. After 72 hours, the CAMs were harvested and examined for changes in the angiogenic index as determined by double blind counting of the number of branch points in the CAM underlying the disk. For kinase assays, the tissue underlying the disk was harvested in RIPA, homogenized with a motorized grinder and Src immunoprecipitated from equivalent amounts of total protein and subjected to an in vitro kinase assay using a FAK-GST fusion protein as a substrate. For the immunofluorescence studies, CAM tissue underlying the disks were frozen in OCT, a cryopreservative, sectioned at 4 µm, fixed in acetone for 1 minute, incubated in 3% normal goat serum for 1 hour, followed by an incubation in primary rabbit anti-phosphorylated ERK antibody as described previously (Eliceiri et al., J. Cell Biol., 140:1255-1263 (1998), washed in PBS and detected with a fluorescent secondary antibody.

A. Activation of Endogenous Src by bFGF or VEGF

To assess the effects of growth factors on Src activity in modulating angiogenesis, the following assays were performed. Tissue extracts of 10 day old chick CAMs that had been exposed to bFGF or VEGF (2 µg/ml) for 2 hours were lysed. Endogenous Src was immunoprecipitated from equivalent amounts of total protein and subjected to an in vitro immune complex kinase assay using a FAK-

GST fusion protein as a substrate, electrophoresed and transferred to nitrocellulose.

The results of the assay are shown in FIG. 5 where the increase in Src activity is evident in the increased density of the gel with either bFGF or VEGF treatment as compared to untreated (mock) samples that are indicative of baseline Src activity in the CAM assay. Both bFGF and VEGF resulted in approximately a 2 fold increase of endogenous Src activity present in the CAM. The above kinase assay blot was also probed with an anti-Src antibody as a loading control for equivalent Src and IgG content.

10        B.     Effect of Retrovirus-Mediated Gene Expression of Src A on Angiogenesis in the Chick CAM

The following assay was performed to assess the effect of mutated Src proteins on angiogenesis in the CAM preparation. For the assay, 9 day old chick CAMs were exposed to RCAS-Src A or RCAS-GFP expressing retroviruses or buffer for 72 hour following the protocol described above.

The results of this assay are shown in FIG. 6A where the level of angiogenesis was quantified as described above. Representative photomicrographs (4x) were taken with a stereomicroscope as shown in FIG. 6B. Baseline endogenous Src activity has an angiogenic index of approximately 50. In contrast, CAMs treated with retroviral vector-expressed RCAS-Src A having a point mutation at amino acid residue position 527 from a tyrosine to a phenylalanine resulted in an enhancement (induction) of angiogenesis of an angiogenic index of approximately 90. The enhancement of Src-A mediated angiogenesis is also evident in the photographs shown in FIG. 6B.

25        C.     Retroviral Expression of Src A Activates Vascular MAP Kinase Phosphorylation

The effect of Src A as compared to growth factors VEGF and PMA on vascular MAP kinase phosphorylation was also assessed following the assay procedures described above and herein. Tissue extracts of 10 day old chick CAMs exposed to VEGF or PMA (another mitogen at a comparable concentration) for 30 minutes were compared to those infected with Src A-expressing retrovirus for 48 hours. Src was then immunoprecipitated from equivalent amounts of total protein extract and subjected to an in vitro immune complex kinase assay using a

FAK-GST fusion protein as a substrate, electrophoresed and transferred to nitrocellulose.

The results of this assay are shown in FIG. 7A where untreated CAMs (NT) exhibit base-line endogenous Src-mediated vascular MAP kinase phosphorylation. Both VEGF and PMA resulted in an approximate 2 fold increase over baseline. In contrast, Src A enhanced the activity approximately 5 to 10 fold over that seen with untreated samples.

Aliquots of the above whole tissue lysates were also measured for endogenous ERK phosphorylation by immunoblotting with an anti-phospho-ERK antibody as shown in FIG. 7B. For this assessment, 10 day old CAMs were infected with either mock RCAS or RCAS that expresses SRC A. After two days, CAMs were dissected, cryopreserved in OCT and sectioned at 4  $\mu$ m. Sections were immunostained with an anti-phosphorylated ERK antibody (New England Biolabs), washed and detected with a goat anti-rabbit FITC-conjugated secondary antibody. Fluorescent images were captured on a cooled-CCD camera (Princeton Inst.). The photomicrographs indicate enhanced immunofluorescence with Src A-treated preparations compared to mock controls.

D. Selective Requirement for Src Activity During VEGF, but Not bFGF-Induced Angiogenesis

To assess the effect of Src modulatory activity on growth factor induced angiogenesis, the following assays were performed. Nine day old chick CAMs were exposed to the retroviral vector preparation that expressed the dominant negative Src mutation referred to as Src 251 or Src K295M as previously described. RCAS-Src 251 or control RCAS-GFP retroviruses or buffer CAMS were treated for 20 hours and then incubated for an additional 72 hours in the presence or absence of bFGF or VEGF.

The level of angiogenesis, quantified as described above, is shown in FIG. 8A. Representative photomicrographs (6x), shown in FIG. 8B, were taken with a stereomicroscope. FIG. 8C illustrates a blot probed with an anti-Src antibody to confirm the expression of Src 251 in transfected cells as compared to mock treatments.

The results of the assays described above indicate that both bFGF and VEGF treated CAMS in the presence of RCAS-GFP controls induced angiogenesis over

the Src-mediated baseline angiogenesis seen with mock or untreated CAM preparations. The expressed dominant negative mutant Src 251 was effective at inhibiting VEGF-induced angiogenesis back to baseline levels while not effective at inhibiting bFGF-mediated angiogenesis. The photomicrographs shown in FIG. 8B pictorially confirm the data shown in FIG. 8A. Thus, retrovirally expressed Src 251 is an effective angiogenesis inhibitor, when angiogenesis is induced with VEGF.

Applications of the Src proteins of this invention with other angiogenesis models as described in the Examples below are contemplated in the present invention.

5     5. Regression of Tumor Tissue Growth With Src Modulators as Measured by In Vivo Rabbit Eye Model Assay

15     The effect of Src modulators on growth factor-induced angiogenesis can be observed in naturally transparent structures as exemplified by the cornea of the eye. New blood vessels grow from the rim of the cornea, which has a rich blood supply, toward the center of the cornea, which normally does not have a blood supply. Stimulators of angiogenesis, such as bFGF, when applied to the cornea induce the growth of new blood vessels from the rim of the cornea. Antagonists of angiogenesis, applied to the cornea, inhibit the growth of new blood vessels from the rim of the cornea. Thus, the cornea undergoes angiogenesis through an invasion of endothelial cells from the rim of the cornea into the tough collagen-packed corneal tissue which is easily visible. The rabbit eye model assay therefore provides an in vivo model for the direct observation of stimulation and inhibition of angiogenesis following the implantation of compounds directly into the cornea of the eye.

25     A.     In Vivo Rabbit Eye Model Assay

1)     Angiogenesis Induced by Growth Factors

Angiogenesis is induced in the in vivo rabbit eye model assay with growth factors bFGF or VEGF and is described in the following sections.

30     a.     Preparation of Hydron Pellets Containing Growth Factor and Monoclonal Antibodies

Hydron polymer pellets containing growth factor are prepared as described by D'Amato, et al., Proc. Natl. Acad. Sci., USA, 91:4082-

4085 (1994). The individual pellets contain 650 ng of the growth factors separately bound to sucralfate (Carafet, Marion Merrell Dow Corporation) to stabilize the growth factor and ensure its slow release into the surrounding tissue. In addition, hydron pellets are prepared containing a desired Src-expressing retrovirus as previously described. The pellets are cast in specially prepared Teflon pegs that have a 2.5 mm core drilled into their surfaces. Approximately 12 ul of casting material is placed into each peg and polymerized overnight in a sterile hood. Pellets are then sterilized by ultraviolet irradiation. Effects of Src proteins are then assessed as previously described.

6. In Vivo Regression of Tumor Tissue Growth With Src Modulators As Measured by Chimeric Mouse:Human Assay

An in vivo chimeric mouse:human model is generated by replacing a portion of skin from a SCID mouse with human neonatal foreskin. The in vivo chimeric mouse:human model is prepared essentially as described in Yan, et al., J. Clin. Invest., 91:986-996 (1993). Briefly, a 2 cm<sup>2</sup> square area of skin is surgically removed from a SCID mouse (6-8 weeks of age) and replaced with a human foreskin. The mouse is anesthetized and the hair removed from a 5 cm<sup>2</sup> area on each side of the lateral abdominal region by shaving. Two circular graft beds of 2 cm<sup>2</sup> are prepared by removing the full thickness of skin down to the fascia. Full thickness human skin grafts of the same size derived from human neonatal foreskin are placed onto the wound beds and sutured into place. The graft is covered with a Band-Aid which is sutured to the skin. Micropore cloth tape is also applied to cover the wound.

The M21-L human melanoma cell line or MDA 23.1 breast carcinoma cell line (ATCC HTB 26;  $\alpha_v\beta_3$  negative by immunoreactivity of tissue sections with mAb LM609), are used to form the solid human tumors on the human skin grafts on the SCID mice. A single cell suspension of  $5 \times 10^6$  M21-L or MDA 23.1 cells is injected intradermally into the human skin graft. The mice are then observed for 2 to 4 weeks to allow growth of measurable human tumors.

After a measurable tumor is established, retrovirus preparations of the present invention or PBS is injected into the mouse tail vein. Following a 2-3 week period, the tumor is excised and analyzed by weight and histology. The effect of expressed Src proteins of the present invention on the tumors is then assessed.



7. In Vitro Regression of Human Tumor Tissue Growth With Src Modulators As Measured by CAM Assay

Tumor growth depends on angiogenesis (Folkman, 1992; Weidner et al., 1991; Brooks et al., 1994b). In fact, recent reports suggest that tumor growth is susceptible to the anti-angiogenic effects of VEGF receptor antagonists (Kim et al., 1993). Therefore, we examined whether suppression of angiogenesis by delivery of kinase-deleted Src 251 would influence the growth of a human medulloblastoma (DAOY), a highly angiogenic tumor known to produce VEGF and very little bFGF (data not shown).

The 3 and 6 day DAOY medulloblastoma tumor growth assays were performed in the chick CAM essentially as previously described (Brooks et al., 1994).  $5 \times 10^6$  DAOY cells cultured in RPMI 1640 containing 10% fetal calf serum were washed and seeded on the CAM of a 10 day embryo to produce DAOY tumor fragments. After 7 days 50 mg tumor fragments were dissected and reseeded on another 10 day embryo and incubated for another 3 or 6 days with the topical application (25 $\mu$ l) of either control RCAS-GFP retrovirus, RCAS-Src 251, or mock treatment. Using the whole tissue confocal imaging of infected tumors as a guide we were able to determine that there was significant expression of the RCAS constructs around and within the tumor fragment with this topical approach. Tumor resections and weighing were performed in a double blind manner removing only the easily definable solid tumor mass (Brooks et al., 1994). The wet tumor weights after 3 or 6 days were compared with initial weight and the percent change of tumor weight determined for each group.

These tumors readily grow on the CAM and produces active angiogenesis (FIG. 9) allowing us to selectively target the avian-derived tumor vasculature by using an avian-specific RCAS retrovirus.

FIG. 9 depicts results that show retroviral delivery of RCAS-Src 251 to human tumors growing on the chick CAM reverses tumor growth. FIG. 9A shows human medulloblastomas that were grown on the CAM of chick embryos as described above. Retrovirus containing RCAS-GFP or RCAS-Src 251 was topically applied to preestablished tumors of greater than 50 mg. A representative micrograph of a medulloblastoma tumor fragment infected with RCAS-GFP expressing GFP reveals exclusive expression in the tumor blood vessels

(arrowhead) as detected by optical sectioning with a Bio Rad laser confocal scanning microscope (bar=500 $\mu$ m). FIG. 9B shows results from tumors treated as above that were allowed to grow for 3 or 6 days after which they were resected and wet weights determined. Data are expressed as the mean change in tumor weight (from the 50 mg tumor starting weight) +/- SEM of 2 replicates. RCAS-Src 251 had a significant impact on tumor growth after 3 days (\*,  $P < 0.002$ ) and 6 days (\*\*,  $P < 0.05$ ). FIG. 9C shows representative stereomicrographs of medulloblastoma tumors surgically removed from the embryos were taken with an Olympus stereomicroscope (bar=350 $\mu$ m). (Lower panel) A high magnification micrograph of each tumor showing the vasculature of each tumor in detail (bar=350 $\mu$ m). The arrowhead indicates blood vessel disruption in RCAS-Src251-treated tumors.

The results show that delivery of RCAS containing Src 251 to preestablished medulloblastomas resulted in selective viral expression in the tumor-associated blood vessels (FIG. 9A) and this ultimately led to the regression of these tumors within the span of six days (FIG. 9B). Importantly, the tumor-associated blood vessels in animals treated with virus containing Src 251 were severely disrupted and fewer in number compared to the tumor vessels in control animals (FIG. 9C). The fact that RCAS-GFP infected tumors showed GFP localization only in the tumor vasculature suggests that the anti-tumor effects observed with retrovirally delivered Src 251 were due to its anti-angiogenic properties.

The foregoing examples and the accompanying description are illustrative, and are not to be taken as limiting. The present invention also is not to be limited in scope by the cell line deposited, since the deposited embodiment is intended as a single illustration of one aspect of the invention. Any cell line that is functionally equivalent is within the scope of this invention. The deposit of material does not constitute an admission that the written description herein contained is inadequate to enable the practice of any aspect of the invention, including the best mode thereof, nor is it to be construed as limiting the scope of the claims to the specific illustration that it represents. Indeed, various modifications of the invention in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description and fall within the scope of the appended claims.

## What Is Claimed Is:

1. An article of manufacture comprising packaging material and a pharmaceutical composition contained within said packaging material, wherein said pharmaceutical composition is capable of modulating angiogenesis in a tissue associated with a disease condition, wherein said packaging material comprises a label which indicates that said pharmaceutical composition can be used for treating disease conditions by modulating angiogenesis, and wherein said pharmaceutical composition comprises a Src protein or an oligonucleotide having a nucleotide sequence capable of expressing said protein.
2. The article of manufacture of claim 1 wherein said Src protein is an active Src protein and said modulating potentiates angiogenesis.
3. The article of manufacture of claim 2 wherein said active Src protein is Src A.
4. The article of manufacture of claim 2 wherein said tissue has poor circulation.
5. The article of manufacture of claim 1 wherein said tyrosine kinase Src protein is an inactive Src protein and said modulating inhibits angiogenesis.
6. The article of manufacture of claim 5 wherein said inactive Src protein is Src 251 or Src K295M.
7. The article of manufacture of claim 5 wherein said tissue is inflamed and said condition is arthritis or rheumatoid arthritis.
8. The article of manufacture of claim 5 wherein said tissue is a solid tumor or solid tumor metastasis.
9. The article of manufacture of claim 8 wherein said administering is conducted in conjunction with chemotherapy.
10. The article of manufacture of claim 5 wherein said tissue is retinal tissue and said condition is retinopathy, diabetic retinopathy or macular degeneration.
11. The article of manufacture of claim 5 wherein said tissue is at the site of coronary angioplasty and said condition is restenosis.
12. The article of manufacture of claim 1 wherein said administering comprises intravenous, transdermal, intrasynovial, intramuscular, or oral administration.

13. The article of manufacture of claim 1 wherein said administering comprises a single dose intravenously.

14. The article of manufacture of claim 1 wherein said pharmaceutical composition further comprises a liposome.

5 15. The article of manufacture of claim 1 wherein said pharmaceutical composition comprises a viral expression vector capable of expressing said nucleotide sequence.

16. The article of manufacture of claim 1 wherein said pharmaceutical composition comprises an non-viral expression vector capable of expressing said  
10 nucleotide sequence.

17. A method for modulating angiogenesis in a tissue associated with a disease condition comprising administering to said tissue a pharmaceutical composition comprising a Src protein or a nucleotide sequence capable of expressing said protein.

15 18. The method of claim 17 wherein said Src protein is an active Src protein and said modulating potentiates angiogenesis.

19. The method of claim 18 wherein said active Src protein is Src A.

20. The method of claim 18 wherein said tissue has poor circulation.

21. The method of claim 17 wherein said Src protein is an inactive Src  
20 protein and said modulating inhibits angiogenesis.

22. The method of claim 21 wherein said inactive Src protein is Src 251 or Src K295M.

23. The method of claim 21 wherein said tissue is inflamed and said condition is arthritis or rheumatoid arthritis.

25 24. The method of claim 21 wherein said tissue is a solid tumor or solid tumor metastasis.

25. The method of claim 24 wherein said administering is conducted in conjunction with chemotherapy.

26. The method of claim 21 wherein said tissue is retinal tissue and said  
30 condition is retinopathy, diabetic retinopathy or macular degeneration.

27. The method of claim 21 wherein said tissue is at the site of coronary angioplasty and said tissue is at risk for restenosis.

28. The method of claim 17 wherein said administering comprises intravenous, transdermal, intrasynovial, intramuscular, or oral administration.

29. The method of claim 17 wherein said administering comprises a single dose intravenously.

5 30. The method of claim 17 wherein said pharmaceutical composition further comprises a liposome.

31. The method of claim 17 wherein said pharmaceutical composition comprises an retroviral expression vector capable of expressing said nucleotide sequence.

10 32. The method of claim 17 wherein said pharmaceutical composition comprises an non-viral expression vector capable of expressing said nucleotide sequence.

33. A pharmaceutical composition for stimulating angiogenesis in a target mammalian tissue comprising a viral gene transfer vector containing a nucleic acid and pharmaceutically acceptable carrier or excipient; said nucleic acid having a nucleic acid segment encoding for a src protein, said src protein having any amino acid residue at codon 527 except for tyrosine, serine or threonine.

15 34. A pharmaceutical composition for stimulating angiogenesis in a target mammalian tissue comprising a non-viral gene transfer vector containing a nucleic acid and pharmaceutically acceptable carrier or excipient; said nucleic acid having a nucleic acid segment encoding for a src protein and said src protein having any amino acid residue at codon 527 except tyrosine, serine or threonine.

35. A pharmaceutical composition for inhibiting angiogenesis in a target mammalian tissue comprising a viral gene transfer vector containing a nucleic acid and pharmaceutically acceptable carrier or excipient; said nucleic acid having a nucleic acid segment encoding for a src protein having no kinase activity.

25 36. A pharmaceutical composition for inhibiting angiogenesis in a target mammalian tissue comprising a non-viral gene transfer vector containing a nucleic acid and pharmaceutically acceptable carrier or excipient; said nucleic acid having a nucleic acid segment encoding for a src protein, said src protein having no kinase activity.

30 37. A pharmaceutical composition for stimulating angiogenesis in a target mammalian tissue comprising a therapeutic amount of a src protein in a

pharmaceutically acceptable carrier or excipient; said src protein having any amino acid residue at codon 527 except tyrosine, serine or threonine.

38. A pharmaceutical composition for inhibiting angiogenesis in a target mammalian tissue comprising a a src protein in a pharmaceutically acceptable carrier or excipient; said src protein having no kinase activity.
- 5

## CHICKEN c-SRC cDNA

(SEQ ID NO:2)

1 tctgacaccc atctgtctgt ctgtctgtgt gctgcaggag ctgagctgac tctgtctgtg  
61 cctcgcgtac cactgtggcc aggcggtagc tgggacgtgc agcccaccac catggggagc  
121 agcaagagca agcccaagga cccagccag cgccggcgca gcctggagcc acccgacagc  
181 acccaccacg ggggattccc agcctcgag accccaaca agacagcagc ccccgacacg  
241 caccgcaccc ccagccgctc ctttgggacc gtggccaccg agcccaagct cttcgggggc  
301 ttcaacactt ctgacaccgt tacgtcgccg cagcgtgccg gggcactggc tggcggcgct  
361 accactttcg tggctctcta cgactacgag tcccggactg aaacggactt gtcctcaag  
421 aaaggagaac gcctgcagat tgtcaacaac acggaagggtg actggtggct ggctcattcc  
481 ctactacag gacagacggg ctacatcccc agtaactatg tcgcgccctc agactccatc  
541 cagggtgaag agtggctact tgggaagatc actcgtcggg agtccgagcg gctgtctgct  
601 aaccccgaag accccggggg aacctcttg gtccgggaga gcgagacgac aaaagggtgcc  
661 tattgcctct ccgtttctga ctttgacaac gccaaggggc tcaatgtgaa gcactacaag  
721 atccgcaagc tggacagcgg cggtctctac atcacctcac gcacacagtt cagcagcctg  
781 cagcagctgg tggcctacta ctccaacat gctgatggct tgtgccaccg cctgaccaac  
841 gtctgcccc cgtccaagcc ccagaccag ggactcgcca aggacgcgtg ggaaatcccc  
901 cgggagtcgc tgcggctgga ggtgaagctg gggcagggct gctttggaga ggtctggatg  
961 gggacctgga acggcaccac cagagtggcc ataaagactc tgaagcccgg caccatgtcc  
1021 ccggaggcct tctgcagga agcccaagt atgaagaagc tccggcatga gaagtgggt  
1081 cagctgtacg cagtgggtgc ggaagagccc atctacatcg tcaatgagta catgagcaag  
1141 gggagcctcc tggatttct gaaggagag atgggcaagt acctgcggct gccacagctc  
1201 gtcgatatgg ctgtcagat tgcacccggc atggcctatg tggagaggat gaactacgtg  
1261 caccgagacc tgcggcgggc caacatcctg gtgggggaga acctggtgtg caagggtggct  
1321 gactttgggc tggcacgcct catcgaggac aacgagtaca cagcacggca aggtgccaag  
1381 ttcccatca agtggacagc ccccgaggca gccctctatg gccggttac catcaagtgc  
1441 gatgtctggt ccttcggcat cctgtgact gagctgacca ccaagggcgg ggtgccatac  
1501 ccagggatgg tcaacaggga ggtgctggac caggtggaga ggggctaccg catgccctgc  
1561 ccgcccagat gcccagatc gctgcatgac ctcatgtgcc agtgcgtggc gagggaccct  
1621 gaggagcggc ccactttga gtacctgcag gccttcctgg aggactactt cacctcgaca  
1681 gagccccagt accagcctgg agagaacct taggcctgga gtcctcctg gaccagaggc  
1741 ctgcgtgtgg ggtacagg

FIG. 1

2 / 10

## CHICKEN cSRC ENCODED PROTEIN

(SEQ ID NO:3)

MGSSKSKPKDPSQRRRSLEPPDSTHHGGFPASQTPNKTAA  
PDTHRTPSRSFSGTVATEPKLFGGFNTSDTVTSPQRAGALA  
GGVTTFVALYDYESRTETDLSFKKGERLQIVNNTGEDWWL  
AHS�TTGQTGYIPSNYVAPSDSIQAEWYFGKITRRESER  
LLNPENPRGTFLVRESETTKGAYCLSVSDFDNAKGLNVK  
HYKIRKLDSGGFYITSRTQFSSLQQLVAYYSKHADGLCHR  
LTNVCPTSKPQTQGLAKDAWEIPRESLRLEVKLGGQCFGE  
VWMGTWNGTTRVAIKTLKPGTMSPEAFLQEAQVMKKLRHE  
KLVQLYAVVSEEPYIVTEYMSKGSLLDFLKGEMGKYLRL  
PQLVDMAAQIASGMAYVERMNYVHRDLRAANILVGENL  
VCKVADFGLARLIEDNEYTARQGAKFPIKWTAPEAALYGR  
FTIKSDVWSFGILLTELTTKGRVPYPGMVNREVLDQVERG  
YRMPCPPECPESLHDLMCQCWRRDPEERPTFEYLQAFLE  
DYFTSTEPQYQPGENL

FIG. 2



3/10

## HUMAN c-SRC cDNA

(SEQ ID NO:4)

```

1  gcgcgcgcgc cgcaggccg tgatgccgc cgcgcggagg tggcccggac cgcagtcccc
61  caagagagct ctaatggtac caagtacag gtggcctta ctgtactcg gggacgccag
121 agtcctgag aagatgtcag caatacaggc cgctggcca tccgtacag aatgtattgc
181 caagtacaac ttccacggca ctgccgagca ggacctgccc ttctgcaaag gagacgtgct
241 caccattgtg gccgtacca aggacccaa ctggtacaaa gccaaaaaca aggtgggccc
301 tgagggcac atcccagcca actacgtcca gaagcgggag ggcgtgaagg cgggtaccaa
361 actcagcctc atgccttgg tccacggcaa gatcacacgg gagcaggctg agcggcttct
421 gtacccggcg gagacaggcc tgtcctggt gcgggagagc accaactacc ccggagacta
481 cacgctgtgc gtgagctgcg acggcaaggt ggagcactac cgcacatgt accatgccag
541 caagctcagc atcgacgagg aggtgtactt tgagaacctc atgcagctgg tggagcacta
601 cacctcagac gcagatggac tctgtacgc cctcattaaa ccaaaggta tggaggggcac
661 agtggcggcc caggatgagt tctaccgag cggctgggcc ctgaacatga aggagctgaa
721 gctgctgcag accatcggga agggggagtt cggagacgtg atgctgggcg attaccgagg
781 gaacaaagtc gccgtcaagt gcattaagaa cgacgccact gccaggcct tcctggctga
841 agcctcagtc atgacgcaac tgcggcatag caacctggtg cagctcctgg gcgtgatcgt
901 ggaggagaag ggcgggctct acatcgtcac tgagtacatg gccaaaggga gccttgtgga
961 ctacctgcgg tctaggggtc ggtagtctg gggcggagac tgtctctca agttctcgt
1021 agatgtctgc gaggccatgg aatacctgga gggcaacaat ttctgtcgc gagacctggc
1081 tgcccgaat gtgtggtgt ctgaggacaa cgtggccaag gtcagcgaact ttggtctac
1141 caaggaggcg tccagcacc aggacacggg caagctgcca gtcaagtga cagccctga
1201 ggccctgaga gagaagaaat tctccactaa gtctgacgtg tggagttcg gaatcctct
1261 ctgggaaatc tactccttg ggcgagtgc ttatccaaga attcccctga aggacgtcgt
1321 ccctcgggtg gagaagggt acaagatgga tgccccgcac ggctgcccgc cgcagtcta
1381 tgaagtcag aagaactgt ggacactgga cgcggccatg cggccctcct tctacagct
1441 ccgagagcag cttgagcaca tcaaaccca cgagctgcac ctgtgacggc tggcctccgc
1501 ctgggtcatg ggcctgtggg gactgaacct ggaagatcat ggacctggtg cccctgtca
1561 ctggggccga gctgaactg agccccagcg ggctggcggg ccttttct gcgtccagc
1621 ctgcaccct ccggccccgt ctctcttga cccacctgtg gggcctgggg agccactga
1681 ggggccaggg aggaaggagg ccacggagcg ggaggcagcg cccaccacg tcgggcttc
1741 ctggcctccc gccactcgcc ttcttagagt ttattcctt tcctttttg agattttt
1801 tccgtgtgtt tttttttat ttttttcaa gataaggaga aagaagtac ccagcaaat
1861 ggcattttac aagaagtacg aatcttatt ttctgtcct gccctgagg gtggggggga
1921 ccgggcccc ctctaggac cctcgcccc agcctcatt cccattctgt gtccatgtc
1981 cgtgtctcc tcggtcgccc cgtgttgcg cttgacctg ttgactgtt tgcacgccc
2041 cgaggcagac gtctgtcagg ggcttgatt tcgtgtgcc ctgccaccg cccaccgcc
2101 ttgtgagatg gaattgtaat aaaccacgcc atgaggacac cgcggccgc ctcggcgctt
2161 cctccaccga aaaaaaaaa aaaaaaa

```

FIG. 3

4 / 10


## HUMAN c-SRC ENCODED PROTEIN

(SEQ ID NO:5)

MSAIQAAWPSGTECIAKYNFHGTAEQDLPFCKGDVLTIVAVTKD  
PNWYKAKNKVGREGIIPANYVQKREGVKAGTKLSLMPWFHGKIT  
REQAERLLYPPETGLFLVRESTNYPGDYTL CVSCDGKVEHYRIMY  
HASKLSIDEEVYFENLMQLVEHYTSDADGLCTRLIKPKVMEGTVA  
AQDEFYRSGWALNMKELKLLQTIGKGEFGDVMLGDYRGNKVAV  
KCIKNATAQAFLAEASVMTQLRHSNLVQLLGVIVEEKGGLYIVTE  
YMAKGSLVDYLRSRGRSVLGGDCLLKFSLDVCEAMEYLEGNFVH  
RDLAARNVLVSEDNVAKVSDFGLTKEASSTQDTGKLPVKWTAPEAL  
REKKFSTKSDVWSFGILLWEIYSFGRVPYPRIPLKD VVPRVEKGYKM  
DAPDGCPPAVYEVMKNCWHLDAAMRPSFLQLREQLEHIKTHELHL



FIG. 4

### Activation of endogenous Src activity by bFGF and VEGF


**FIG. 5**

6/10


Effect of RCAS-mediated expression of  
Src A on angiogenesis in the chick CAM

**FIG. 6A****FIG. 6B****SUBSTITUTE SHEET (RULE 26)**

# Retroviral expression of Src A activates vascular MAP kinase phosphorylation



Mock



Src A







FIG. 7

SUBSTITUTE SHEET (RULE 26)

8/10

# Selective requirement for Src activity during VEGF, but not bFGF-induced angiogenesis

**FIG. 8A****FIG. 8B****FIG. 8C**

9 / 10




FIG. 9A




FIG. 9B




FIG. 9C

SUBSTITUTE SHEET (RULE 26)

10/10




FIG. 10



1/18

## SEQUENCE LISTING

<110> The Scripps Research Institute et al.

<120> METHODS AND COMPOSITIONS USEFUL FOR MODULATION OF  
ANGIOGENESIS USING TYROSINE KINASE SRC

<130> TSRI 651.1

<140> Not yet known  
<141> To be determined

<150> 60/087,220  
<151> 1998-05-29

<160> 6

<170> PatentIn Ver. 2.0

<210> 1  
<211> 11627  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:RCASBP(A) based  
on avian sarcoma virus

<220>  
<221> misc\_feature  
<222> (7649)..(11258)  
<223> pBR322 sequences

<220>  
<221> LTR  
<222> (7166)..(7494)  
<223> upstream

<220>  
<221> LTR  
<222> (1)..(101)  
<223> upstream (numbering begins at the upstream R)

<220>  
<221> misc\_feature  
<222> (11394)..(11623)  
<223> U3

<220>  
<221> misc\_feature  
<222> (1)..(21)  
<223> R

<220>  
<221> misc\_feature  
<222> (22)..(101)  
<223> U5

<220>  
<221> misc\_feature  
<222> (102)..(119)

<220>  
<221> LTR

2/18

<222> (7166)..(7494)  
<223> downstream

<220>  
<221> misc\_feature  
<222> (7166)..(7393)  
<223> U3

<220>  
<221> misc\_feature  
<222> (7394)..(7414)  
<223> R

<220>  
<221> misc\_feature  
<222> (7415)..(7494)  
<223> U5

<220>  
<221> misc\_feature  
<222> (7154)..(7165)  
<223> PPT

<220>  
<221> misc\_feature  
<222> (388)..(391)  
<223> splice donor (AGGT)

<220>  
<221> misc\_feature  
<222> (5074)..(5077)  
<223> env splice acceptor (AGGC)

<220>  
<221> misc\_feature  
<222> (6982)..(6985)  
<223> ClaI splice acceptor (AGGA)

<220>  
<221> gene  
<222> (372)..(902)  
<223> gag p19

<220>  
<221> gene  
<222> (909)..(1094)  
<223> gag p10

<220>  
<221> gene  
<222> (1095)..(1814)  
<223> gag p27

<220>  
<221> gene  
<222> (1843)..(2108)  
<223> gag p12

<220>  
<221> gene  
<222> (2109)..(2480)  
<223> gag p15

<220>

3/18

<221> misc\_signal  
<222> (2481)..(2483)  
<223> gag stop

<220>  
<221> gene  
<222> (2501)..(4216)  
<223> pol RT

<220>  
<221> gene  
<222> (4217)..(5185)  
<223> pol IN

<220>  
<221> misc\_signal  
<222> (5186)..(5188)  
<223> pol stop

<220>  
<221> gene  
<222> (5244)..(6263)  
<223> env gp85

<220>  
<221> gene  
<222> (6264)..(6878)  
<223> env gp37

<220>  
<221> misc\_signal  
<222> (6879)..(6881)  
<223> env stop

<220>  
<221> misc\_feature  
<222> (7027)  
<223> ClaI site/ the ClaI site in gag is methylated in  
Dam+ strains and does not cut.

<400> 1  
gccatttgac cattcaccac attggtgtgc acctgggttg atggccggac cgttgattcc 60  
ctgacgacta cgagcacctg catgaagcag aaggcttcat ttggtgacct cgacgtgata 120  
gttaggggaat agtggtcggc cacagacggc gtggcgatcc tgtctccatc cgtctcgtct 180  
atcgggaggg gagttcgatg accctgggtg agggggctgc ggcttaggga ggcagaagct 240  
gagtaccgtc ggaggagct ccaggggccc gagcgactga cccctgccga gaactcagag 300  
ggtcgtcggg agacggagag tgagcccgc gaccaccca ggcacgtctt tggtcggcct 360  
gcggatcaag catggaagcc gtcattaagg tgatttcgtc cgcggtgtaa acctattgcg 420  
ggaaaatctc tccttctaag aaggaaatcg gggccatgtt gtccctgtta caaaaggaag 480  
ggttgcttat gtctccctca gatttatatt ctccggggtc ctgggatccc atcactgcgg 540  
cgctctccca gcgggcaatg gtacttgga aatcgggaga gttaaaaacc tggggattgg 600  
ttttgggggc attgaaggcg gtcgagagg aacagggttac atctgagcaa gcaaagtttt 660  
ggttgggatt agggggaggg aggggtcttc cccagggtcc ggagtgcac gagaaaccag 720

4/18

ctacggagcg gogaatcgac aaaggggagg aggtgggaga aacaactgtg cagcgagatg 780  
cgaagatggc gccagaggaa gcggccacac ctaaaaccgt tggcacatcc tgctatcatt 840  
gcggaacagc tgttggctgc aattgcgcca ccgccacagc ctcgccccct cctccccctt 900  
atgtggggag tggtttgtat ccttccctgg cgggggtggg agagcagcag ggccaggag 960  
ataacacgtc tcggggggcg gagcagccaa gggaggagcc agggcacgag ggtcaggccc 1020  
ctggggccggc cctgactgac tgggcaaggg taaggaggga gcttgcgagt actggtccgc 1080  
ccgtgggtggc catgcctgta gtgattaaga cagagggacc cgcctggacc cctctggagc 1140  
caaaattgat cacaagactg gctgatacgg tcaggaccaa gggcttacga tccccgatca 1200  
ctatggcaga agtgggaagc ctcatgtcct ccccgttgct gccgcatgac gtcacgaatc 1260  
taatgagagt gatttttagga cctgccccat atgccttatg gatggacgct tggggagtcc 1320  
aactccagac gggttatagc gcagccactc gcgacccccg acaccagcgc aacgggtcaag 1380  
ggcgggggga acggactaac ttggatcgat taaagggtct agctgatggg atgggtgggca 1440  
accacaggg tcaggccgca ttattaagac cgggggaatt ggttgctatt acggcgtcgg 1500  
ctctccaggc gtttagagaa gttgcccggc tggcggaacc tgcaggtcca tgggaggaca 1560  
tcacgcaggg accatctgag tcctttgttg attttgccaa tcggcttata aaggcggttg 1620  
aggggtcaga tctccgcct tccgcgcggg ctccggtgat cattgactgc tttaggcaga 1680  
agtcacagcc agatattcag cagcttatac gggcagcacc ctccacgctg accacccag 1740  
gagagataat caaatatgtg ctagacaggc agaagattgc ccctcttacg gatcaaggca 1800  
tagccgcggc catgtcgtct gctatccagc ccttagttat ggcagtagtc aatagagaga 1860  
gggatggaca aactgggtcg ggtggtcgtg cccgagggtc ctgctacact tgtggatccc 1920  
cgggacatta tcaggcacag tgcccgaata aacgaaagtc aggaaacagc cgtgagcgat 1980  
gtcagctgtg tgacgggatg ggacacaacg ctaaacagtg taggaagcgg gatggcaacc 2040  
agggccaacg cccaggaaga ggtctctctt cggggccgtg gcccggccct gagcagcctg 2100  
ccgtctcggt agcgatgaca atggaacata aagatcgccc cttgggttagg gtcattctga 2160  
ctaactctgg gagtcatcca gtcaaacaac gttcgggtgta tatcaccgag ctggttgact 2220  
ccggagcggg catcactatt atttcggagg aggattggcc tactgattgg ccggtgggtg 2280  
acaccgcgaa cccacagatc catggcatag gagggggaat tcccatgcga aaatccccgg 2340  
atatgataga ggtgggggtt attaaccgag acgggtcgtt ggagcgaccc ctgctcctct 2400  
tccccgcagt cgctatgggt agagggagta tcctaggaag agattgtctg cagggcctag 2460  
ggctccgctt gacaaattta tagggagggc cactgttctc actgttgccg tacatctggc 2520  
tattccgctc aaatggaagc cagaccgcac gcctgtgtgg attgaccagt ggccccctcc 2580  
tgaaggtaaa cttgtagggc taacgcaatt agtggaaaaa gaattacagt taggacatat 2640

agagccctca cttagttggt ggaacacacc tgtttttcgt gatccggaag gcttccgggt 2700  
cttatcgctt attgcatgat ttgcgcgctg ttaacgcaa gcttgccct tttggggccg 2760  
tccaacaggg ggcgccagtt ctctccgcgc tcccgctgg ctggccctg atggctctag 2820  
acctcaagga ttgcttcttt tctatccctc ttgcggaaca agatcgcgaa gcttttgcat 2880  
ttacgctccc ctctgtgaat aaccaggccc ccgctcgaag attccaatgg aaggctcttg 2940  
ccaagggat gacctgttct ccactatct gtcagttggt agtgggtcag gtgctcgagc 3000  
cettgcgact caagcaccca gctctgcgca tggtgcatta tatggacgat cttttgctag 3060  
ccgcctcaag tcatgatggg ttggaagcgg cagggaagga ggttatcggg acattggaaa 3120  
gagccgggtt cactatttcg ccggataaga tccagaggga gcccgagta caatatcttg 3180  
ggtacaagtt aggcagtacg tatgtagcac ccgtaggctt ggtagcagaa cccaggatag 3240  
ccaccttggt ggatgttcaa aagctggtgg ggtcacttca gtggcttcgc ccagcgtag 3300  
ggatcccgcc acgactgatg ggtccctttt atgagcagtt acgagggtca gatcctaacg 3360  
aggcgaggga atggaatcta gacatgaaaa tggcctggag agagatcgta cagcttagca 3420  
ctactgctgc cttggaacga tgggaccctg ccagcctct ggaaggagcg gtcgctagat 3480  
gtgaacaggg ggcaataggg gtcctgggac agggactgtc cacacacca aggccatggt 3540  
tgtggttatt ctccaccaa cccaccaagg cgtttactgc ttggttagaa gtgctcacc 3600  
ttttgattac taagctacgc gtttcggcag tgcgaacctt tggcaaggag gttgatatcc 3660  
tctgttgcc tgcagtcttc cgggaggacc ttccgctccc ggaggggatc ctgttagcac 3720  
ttaggggggt tgcaggaaaa atcaggagta gtgacacgcc atctatttt gacattgcgc 3780  
gtccactgca tgtttctctg aaagtgagg ttaccgacca ccctgtgcg ggacccactg 3840  
tctttaccga cgcctcctca agcaccata aaggggtggt agtctggagg gagggccaa 3900  
ggtgggagat aaaagaaata gttgatttgg gggcaagtgt acaacaactg gaggcacgcg 3960  
ctgtggccat ggcacttctg ctgtggccga caacgcccac taatgtagtg actgactctg 4020  
cgtttggtgc gaaaatgtta ctcaagatgg gacaggagg agtcccgtct acagcgcg 4080  
cttttatttt agaggatgcg ttaagccaaa ggtcagccat ggccgctgt ctccacgtgc 4140  
ggagtcattc tgaagtgcc gggtttttca cagaaggaaa tgacgtggca gatagccaag 4200  
ccacctttca agcgtatccc ttgagagagg ctaaagatct tcataccgct ctccatattg 4260  
gaccccgcg gctatccaaa gcgtgtaata tatctatgca gcaggctagg gaggttggtc 4320  
agacctgcc gcattgtaat tcagccctg cgttgaggc cggggtaaac cctaggggtt 4380  
tgggacccct acagatatgg cagacagact ttacgcttga gcctagaatg gctccccgtt 4440  
cctggctcgc tgttactgtg gacaccgct catcagcgat agtcgtaact cagcatggcc 4500  
gtgttacatc ggttgctgca caacatcatt gggccacggc tatcgccgtt ttgggaagac 4560

caaaggccat aaaaacagat aacgggtcct gcttcacgtc cagatccacg cgagagtggc 4620  
tcgcgagatg ggggatagca cacaccaccg ggattccggg aaattcccag ggtcaagcta 4680  
tggtagagcg ggccaaccg ctcctgaaag ataagatccg tgtgctcgcg gagggggacg 4740  
gctttatgaa aagaatcccc accagcaaac aggggggaact attagccaag gcaatgtatg 4800  
ccctcaatca ctttgagcgt ggtgaaaaca caaaaacacc gatacaaaaa cactggagac 4860  
ctaccgttct tacagaagga cccccggta aaatacgaat agagacaggg gagtgggaaa 4920  
aaggatggaa cgtgctggtc tggggacgag gttatgccgc tgtgaaaaac agggacactg 4980  
ataaggttat ttgggtaccc tctcggaaag ttaaaccgga tgtcacccaa aaggatgagg 5040  
tgactaagaa agatgaggcg agccctcttt ttgcaggcat ttctgactgg ataccctggg 5100  
aagacgagca agaaggactc caaggagaaa ccgctagcaa caagcaagaa agacccgagg 5160  
aagacacctt tgctgccaac gagagttaat tatattctca ttattggtgt cctggtcttg 5220  
tgtgaggtta cgggggtaag agctgatgtc cacttactcg agcagccagg gaaccttttg 5280  
attacatggg ccaaccgtac aggccaaacg gatttttgcc tctctacaca gtcagccacc 5340  
tccccctttc aaacatgttt gataggtatc ccgtccccta tttccgaggg tgattttaag 5400  
ggatatgttt ctgatacaaa ttgcaccacc ttgggaactg atcggttagt ctcgtcagcc 5460  
gactttactg gcggaacctga caacagtacc accctcactt atcggaaggt ctcatgcttg 5520  
ttgttaaagc tgaatgtctc tatgtgggat gagccacctg aactacagct gttaggttcc 5580  
cagtctctcc ctaacattac taatattgct cagatttccg gtataaccgg gggatgcgta 5640  
ggcttcagac cacaaggggt tccttggtat ctaggttggt ctagacagga ggccacgcg 5700  
tttctcctta gacaccctc tttctctaaa tccacggaac cgtttacagt ggtgacagcg 5760  
gataggcaca atctttttat ggggagttag tactgcggtg catatggcta cagatttttg 5820  
aacatgtata actgctcaca ggtggggcgg cagtaccgct gtggtaatgc gcgcacgccc 5880  
cgcacgggtc ttctgaaat ccagtgtaca aggagaggag gcaaattgggt taatcaatca 5940  
caggaaatta atgagtcgga gccgttcagc tttacgggtga actgtacagc tagtagtttg 6000  
ggtaatgcc a gtgggtggtg cggaaaagca ggcacgattc tcccgggaaa gtgggtcgac 6060  
agcacacaag gtagtttcac caaaccaaaa gcgctaccac ccgcaatttt cctcattttg 6120  
ggggatcgcg catggcaagg aattcccagt cgtccggtag ggggcccctg ctattttaggc 6180  
aagcttacca tgtagcacc taagcataca gatattctca aggtgcttgt caattcatcg 6240  
cggacaggta taagacgtaa acgaagcacc tcacacctgg atgatacatg ctcatatgaa 6300  
gtgcagcttt ggggtcctac agcaagaatc tttgcatcta tcctagcccc gggggtagca 6360  
gctgcgcaag ccttaagaga aattgagaga ctacgctgtt ggtecggtta acaggctaac 6420  
ttgacaacat cactcctcgg ggacttattg gatgatgtca cgagtattcg acacgcggtc 6480

7/18

ctgcagaacc gagcggctat tgacttcttg ctccctagctc acggccatgg ctgtgaggac 6540  
gttgccggaa tgtgctgttt caatttgagt gatcagagtg agtctataca gaagaagttc 6600  
cagctaataga aggaacatgt caataagatc ggcgtaggata gcgacctaat tgggaagttgg 6660  
ctgcgaggac tattcggggg aataggagaa tgggcccgttc atttgctgaa aggactgctt 6720  
ttggggcttg tagttatattt gttgctagta gtgtgcctgc cttgcctttt gcaaagtta 6780  
tgcggttaata ggagaaagat gattaataac tccatcagct accacacgga atataagaag 6840  
ctgcaaaagg cctgtgggca gcctgaaagc agaatagtat aaggcagtac atgggtggtg 6900  
gtatagcgct tgcgagtgca tcgagcaagg caggaaagac agctattggt aattgtgaaa 6960  
tacgcttttg tctgtgtgct gcaggagctg agctgactct gctggtggcc tcgcgtacca 7020  
ctgtggcatc gatgcgatgt acggggccaga tatacgcgta tctgagggga ctagggtgtg 7080  
tttagcgcaa aagcggggct tcggttgtag gcggttagga gtccccttag gatatagtag 7140  
tttcgctttt gcatagggag ggggaaatgt agtcttatgc aatactcttg tagtcttgca 7200  
acatggtaac gatgagttag caacatgcct tacaaggaga gaaaaagcac cgtgcatgcc 7260  
gattggtgga agtaaggtgg tacgatcgtg ccttattagg aaggcaacag acgggtctga 7320  
catggattgg acgaaccact gaattccgca ttgcagagat attgtattta agtgcctagc 7380  
tcgatacaat aaacgccatt tgaccattca ccacattggt gtgcacctgg gttgatggcc 7440  
ggaccgttga ttccctgacg actacgagca cctgcatgaa gcagaaggct tcatttggtg 7500  
accccgacgt gatagttagg gaatagtggc cggccacaga cggcgtggcg atcctgtctc 7560  
catccgtctc gtctatcggg aggcgacttc gatgacctg gtggaggggg ctgcggctta 7620  
gggaggcaga agctgagtag cgtcggaggg gatccacagg acgggtgtgg tcgccatgat 7680  
cgcgtagtcg atagtggctc caagtagcga agcgagcagg actgggcggc ggccaaagcg 7740  
gtcggacagt gctccgagaa cgggtgcgca tagaaattgc atcaacgcat atagcgctag 7800  
cagcacgcca tagtgactgg cgatgctgtc ggaatggacg atatcccgca agaggcccgg 7860  
cagtaccggc ataaccaagc ctatgcctac agcatccagg gtgacggtgc cgaggatgac 7920  
gatgagcgca ttgtagatt tcatacacgg tgccctgactg cgtagcaat ttaactgtga 7980  
taaactaccg cattaaagct ccaaacttg cgttttctg tgtgaaattg ttatccgctc 8040  
acaattccac acattatacg agccggaagc ataaagtgtg aaacctgggg tgccaatga 8100  
gtgagaattc ttgaagacga aagggcctcg tgatacgctc atttttatag gttaatgtca 8160  
tgataataat ggtttcttag acgtcagggtg gcacttttcg gggaaatgtg cgcggaaccc 8220  
ctatttgttt atttttctaa atacattcaa atatgtatcc gctcatgaga caataaccct 8280  
gataaatgct tcaataatat tgaaaaagga agagtatgag tattcaacat ttccgtgtcg 8340  
cccttattcc cttttttgcg gcattttgcc ttccgtgttt tgctcaccca gaaacgctgg 8400

tgaaagtaaa agatgctgaa gatcagttgg gtgcacgagt gggttacatc gaactggatc 8460  
tcaacagcgg taagatcctt gagagttttc gccccgaaga acgttttcca atgatgagca 8520  
cttttaaaagt tctgctatgt ggcgcggtat tatcccgtgt tgacgcgggg caagagcaac 8580  
tcggtcgccc catacactat tctcagaatg acttggttga gtactcacca gtcacagaaa 8640  
agcatcttac ggatggcatg acagtaagag aattatgcag tgctgccata accatgagtg 8700  
ataaactgac ggccaactta cttctgacaa cgatcggagg accgaaggag ctaaccgctt 8760  
ttttgcacaa catgggggat catgtaactc gccttgatcg ttgggaaccg gagctgaatg 8820  
aagccatacc aaacgacgag cgtgacacca cgatgcctgc agcaatggca acaacgttgc 8880  
gcaaaactatt aactggcgaa ctacttactc tagcttcccg gcaacaatta atagactgga 8940  
tgaggcgga taaagttgca ggaccacttc tgcgctcggc ccttcgggct ggctggttta 9000  
ttgctgataa atctggagcc ggtgagcgtg ggtctcgcgg tatcattgca gcaactgggc 9060  
cagatggtaa gccctcccgt atcgtagtta tctacacgac ggggagtcag gcaactatgg 9120  
atgaacgaaa tagacagatc gctgagatag gtgcctcact gattaagcat tggtaactgt 9180  
cagaccaagt ttactcatat atactttaga ttgatttaaa acttcatttt taattttaaaa 9240  
ggatctaggt gaagatcctt tttgataatc tcatgaccaa aatcccttaa cgtgagtttt 9300  
cgttccactg agcgtcagac cccgtagaaa agatcaaagg atcttcttga gatccttttt 9360  
ttctgcgcgt aatctgctgc ttgcaaacaa aaaaaccacc gctaccagcg gtggtttgtt 9420  
tgccggatca agagctacca actctttttc cgaaggtaac tggttcagc agagcgcaga 9480  
taccaaatac tgtccttcta gtgtagccgt agttaggcca ccacttcaag aactctgtag 9540  
caccgcctac atacctcgt ctgctaatec tgttaccagt ggctgctgcc agtggcgata 9600  
agtcgtgtct taccggggtg gactcaagac gatagttacc ggataaggcg cagcggtcgg 9660  
gctgaacggg gggttcgtgc acacagccca gcttgagcg aacgacctac accgaactga 9720  
gatacctaca gcgtgagcta tgagaaagcg ccacgcttcc cgaagggaga aaggcggaca 9780  
ggtatccggt aagcggcagg gtcggaacag gagagcgcac gaggagctt ccagggggaa 9840  
acgcctggta tctttatagt cctgtcgggt ttcgccacct ctgacttgag cgtcgatttt 9900  
tgtgatgctc gtcagggggg cggagcctat ggaaaaacgc cagcaacgcg gcctttttac 9960  
ggttcctggc cttttgctgg cttttgctc acatgttctt tctgctgta tcccctgatt 10020  
ctgtggataa ccgtattacc gcctttgagt gagctgatac cgctcgccgc agccgaacga 10080  
ccgagcgcag cgagtcagt agcgaggaag cggaagagcg cctgatgcgg tattttctcc 10140  
ttacgcatct gtgcggtatt tcacaccgca tatggtgcac tctcagtaca atctgctctg 10200  
atgccgcata gttaagccag tatacactcc gctatcgcta cgtgactggg tcatggctgc 10260  
gccccgacac ccgccaacac ccgctgacgc gccctgacgg gcttgtctgc tcccggcatc 10320



9/18

```

cgcttacaga caagctgtga cegtctccgg gagctgcatg tgtcagaggt ttccaccgtc 10380
atcaccgaaa cgcgcgagggc agctgcggta aagctcatca gcgtgggtcgt gaagcgattc 10440
acagatgtct gcctgttcat ccgcgtccag ctcggtgagt ttctccagaa gcgttaaatgt 10500
ctggcttctg ataaagcggg ccatgttaag ggcggttttt tcctgtttgg tcaacttgatg 10560
cctccgtgta agggggaatt tctgttcatg ggggtaatga taccgatgaa acgagagagg 10620
atgctcacga tacgggttac tgatgatgaa catgcccggg tactggaacg ttgtgagggt 10680
aaacaactgg cggatatggat gcggcggggac cagagaaaaa tcactcaggg tcaatgccag 10740
cgcttcgtta atacagatgt aggtgttcca cagggtagcc agcagcatcc tgcgatgcag 10800
atccggaaca taatggtgca gggcgtgac ttccgcgttt ccagacttta cgaaacacgg 10860
aaaccgaaga ccattcatgt tgttgctcag gtcgcagacg ttttgcagca gcagtcgctt 10920
cacgttcgct cgcgtatcgg tgattcattc tgctaaccag taaggcaacc ccgccagcct 10980
agccgggtcc tcaacgacag gagcacgatc atgagcacc gtggccagga cccaacgctg 11040
cccgagatgc gccgcgtgcg gctgctggag atggcggacg cgatggatat gttctgcca 11100
gggttggttt gcgcattcac agttctccgc aagaattgat tggctccaat tcttgagtg 11160
gtgaatccgt tagcgaggtg ccgccggctt ccattcaggt cgaggtggcc cggtccatg 11220
caccgcgacg caacgcgggg aggcagacaa ggtatagggc ggcgatgcga tgtacgggcc 11280
agatatacgc gtatctgagg ggactagggg gtgttttagg gaaaagcggg gcttcgggtg 11340
tacgcgggta ggagtccct taggatatag tagtttcgct tttgcatagg gagggggaaa 11400
tgtagtctta tgcaatactc ttgtagtctt gcaacatggg aacgatgagt tagcaacatg 11460
ccttacaagg agagaaaaag caccgtgcat gccgattggg ggaagtaagg tggtagcatc 11520
gtgccttatt aggaaggcaa cagacgggtc tgacatggat tggacgaacc actgaattcc 11580
gcattgcaga gatattgtat ttaagtgcct agctcgatac aataaac 11627

```

```

<210> 2
<211> 1759
<212> DNA
<213> Chicken

```

```

<220>
<221> gene
<222> (1) .. (1759)
<223> chicken c-SRC cDNA

```

```

<220>
<221> CDS
<222> (112) .. (1710)

```

```

<400> 2
tctgacaccc atctgtctgt ctgtctgtgt gctgcaggag ctgagctgac tctgctgtgg 60
cctcgcgtac cactgtggcc aggcggtagc tgggacgtgc agcccaccac c atg ggg 117

```

Met Gly  
1

INSDOCID: <WO\_\_\_\_\_ 9961590A1 |A>

11/18

Asn	Val	Cys	Pro	Thr	Ser	Lys	Pro	Gln	Thr	Gln	Gly	Leu	Ala	Lys	Asp	
		245					250					255				
gcg	tgg	gaa	atc	ccc	cgg	gag	tcg	ctg	cgg	ctg	gag	gtg	aag	ctg	ggg	933
Ala	Trp	Glu	Ile	Pro	Arg	Glu	Ser	Leu	Arg	Leu	Glu	Val	Lys	Leu	Gly	
	260					265					270					
cag	ggc	tgc	ttt	gga	gag	gtc	tgg	atg	ggg	acc	tgg	aac	ggc	acc	acc	981
Gln	Gly	Cys	Phe	Gly	Glu	Val	Trp	Met	Gly	Thr	Trp	Asn	Gly	Thr	Thr	
275					280					285					290	
aga	gtg	gcc	ata	aag	act	ctg	aag	ccc	ggc	acc	atg	tcc	ccg	gag	gcc	1029
Arg	Val	Ala	Ile	Lys	Thr	Leu	Lys	Pro	Gly	Thr	Met	Ser	Pro	Glu	Ala	
				295					300					305		
ttc	ctg	cag	gaa	gcc	caa	gtg	atg	aag	aag	ctc	cgg	cat	gag	aag	ctg	1077
Phe	Leu	Gln	Glu	Ala	Gln	Val	Met	Lys	Lys	Leu	Arg	His	Glu	Lys	Leu	
			310					315					320			
gtt	cag	ctg	tac	gca	gtg	gtg	tcg	gaa	gag	ccc	atc	tac	atc	gtc	act	1125
Val	Gln	Leu	Tyr	Ala	Val	Val	Ser	Glu	Glu	Pro	Ile	Tyr	Ile	Val	Thr	
		325					330					335				
gag	tac	atg	agc	aag	ggg	agc	ctc	ctg	gat	ttc	ctg	aag	gga	gag	atg	1173
Glu	Tyr	Met	Ser	Lys	Gly	Ser	Leu	Leu	Asp	Phe	Leu	Lys	Gly	Glu	Met	
	340					345					350					
ggc	aag	tac	ctg	cgg	ctg	cca	cag	ctc	gtc	gat	atg	gct	gct	cag	att	1221
Gly	Lys	Tyr	Leu	Arg	Leu	Pro	Gln	Leu	Val	Asp	Met	Ala	Ala	Gln	Ile	
355					360					365					370	
gca	tcc	ggc	atg	gcc	tat	gtg	gag	agg	atg	aac	tac	gtg	cac	cga	gac	1269
Ala	Ser	Gly	Met	Ala	Tyr	Val	Glu	Arg	Met	Asn	Tyr	Val	His	Arg	Asp	
				375					380					385		
ctg	cgg	gcg	gcc	aac	atc	ctg	gtg	ggg	gag	aac	ctg	gtg	tgc	aag	gtg	1317
Leu	Arg	Ala	Ala	Asn	Ile	Leu	Val	Gly	Glu	Asn	Leu	Val	Cys	Lys	Val	
			390					395					400			
gct	gac	ttt	ggg	ctg	gca	cgc	ctc	atc	gag	gac	aac	gag	tac	aca	gca	1365
Ala	Asp	Phe	Gly	Leu	Ala	Arg	Leu	Ile	Glu	Asp	Asn	Glu	Tyr	Thr	Ala	
		405					410					415				
cgg	caa	ggt	gcc	aag	ttc	ccc	atc	aag	tgg	aca	gcc	ccc	gag	gca	gcc	1413
Arg	Gln	Gly	Ala	Lys	Phe	Pro	Ile	Lys	Trp	Thr	Ala	Pro	Glu	Ala	Ala	
	420					425					430					
ctc	tat	ggc	cgg	ttc	acc	atc	aag	tcg	gat	gtc	tgg	tcc	ttc	ggc	atc	1461
Leu	Tyr	Gly	Arg	Phe	Thr	Ile	Lys	Ser	Asp	Val	Trp	Ser	Phe	Gly	Ile	
435					440					445					450	
ctg	ctg	act	gag	ctg	acc	acc	aag	ggc	cgg	gtg	cca	tac	cca	ggg	atg	1509
Leu	Leu	Thr	Glu	Leu	Thr	Thr	Lys	Gly	Arg	Val	Pro	Tyr	Pro	Gly	Met	
				455					460					465		
gtc	aac	agg	gag	gtg	ctg	gac	cag	gtg	gag	agg	ggc	tac	cgc	atg	ccc	1557
Val	Asn	Arg	Glu	Val	Leu	Asp	Gln	Val	Glu	Arg	Gly	Tyr	Arg	Met	Pro	
			470					475					480			
tgc	ccg	ccc	gag	tgc	ccc	gag	tcg	ctg	cat	gac	ctc	atg	tgc	cag	tgc	1605
Cys	Pro	Pro	Glu	Cys	Pro	Glu	Ser	Leu	His	Asp	Leu	Met	Cys	Gln	Cys	
		485					490					495				
tgg	cgg	agg	gac	cct	gag	gag	cgg	ccc	act	ttt	gag	tac	ctg	cag	gcc	1653

12/18

Trp Arg Arg Asp Pro Glu Glu Arg Pro Thr Phe Glu Tyr Leu Gln Ala  
 500 505 510

ttc ctg gag gac tac ttc acc tcg aca gag ccc cag tac cag cct gga 1701  
 Phe Leu Glu Asp Tyr Phe Thr Ser Thr Glu Pro Gln Tyr Gln Pro Gly  
 515 520 525 530

gag aac cta taggcctgga gctcctcctg gaccagaggg ctcgctgtgg ggtacaggg 1759  
 Glu Asn Leu

<210> 3  
 <211> 533  
 <212> PRT  
 <213> Chicken

<400> 3  
 Met Gly Ser Ser Lys Ser Lys Pro Lys Asp Pro Ser Gln Arg Arg Arg  
 1 5 10 15  
 Ser Leu Glu Pro Pro Asp Ser Thr His His Gly Gly Phe Pro Ala Ser  
 20 25 30  
 Gln Thr Pro Asn Lys Thr Ala Ala Pro Asp Thr His Arg Thr Pro Ser  
 35 40 45  
 Arg Ser Phe Gly Thr Val Ala Thr Glu Pro Lys Leu Phe Gly Gly Phe  
 50 55 60  
 Asn Thr Ser Asp Thr Val Thr Ser Pro Gln Arg Ala Gly Ala Leu Ala  
 65 70 75 80  
 Gly Gly Val Thr Thr Phe Val Ala Leu Tyr Asp Tyr Glu Ser Arg Thr  
 85 90 95  
 Glu Thr Asp Leu Ser Phe Lys Lys Gly Glu Arg Leu Gln Ile Val Asn  
 100 105 110  
 Asn Thr Glu Gly Asp Trp Trp Leu Ala His Ser Leu Thr Thr Gly Gln  
 115 120 125  
 Thr Gly Tyr Ile Pro Ser Asn Tyr Val Ala Pro Ser Asp Ser Ile Gln  
 130 135 140  
 Ala Glu Glu Trp Tyr Phe Gly Lys Ile Thr Arg Arg Glu Ser Glu Arg  
 145 150 155 160  
 Leu Leu Leu Asn Pro Glu Asn Pro Arg Gly Thr Phe Leu Val Arg Glu  
 165 170 175  
 Ser Glu Thr Thr Lys Gly Ala Tyr Cys Leu Ser Val Ser Asp Phe Asp  
 180 185 190  
 Asn Ala Lys Gly Leu Asn Val Lys His Tyr Lys Ile Arg Lys Leu Asp  
 195 200 205  
 Ser Gly Gly Phe Tyr Ile Thr Ser Arg Thr Gln Phe Ser Ser Leu Gln  
 210 215 220  
 Gln Leu Val Ala Tyr Tyr Ser Lys His Ala Asp Gly Leu Cys His Arg  
 225 230 235 240  
 Leu Thr Asn Val Cys Pro Thr Ser Lys Pro Gln Thr Gln Gly Leu Ala  
 245 250 255

13/18

Lys Asp Ala Trp Glu Ile Pro Arg Glu Ser Leu Arg Leu Glu Val Lys  
 260 265 270  
 Leu Gly Gln Gly Cys Phe Gly Glu Val Trp Met Gly Thr Trp Asn Gly  
 275 280 285  
 Thr Thr Arg Val Ala Ile Lys Thr Leu Lys Pro Gly Thr Met Ser Pro  
 290 295 300  
 Glu Ala Phe Leu Gln Glu Ala Gln Val Met Lys Lys Leu Arg His Glu  
 305 310 315 320  
 Lys Leu Val Gln Leu Tyr Ala Val Val Ser Glu Glu Pro Ile Tyr Ile  
 325 330 335  
 Val Thr Glu Tyr Met Ser Lys Gly Ser Leu Leu Asp Phe Leu Lys Gly  
 340 345 350  
 Glu Met Gly Lys Tyr Leu Arg Leu Pro Gln Leu Val Asp Met Ala Ala  
 355 360 365  
 Gln Ile Ala Ser Gly Met Ala Tyr Val Glu Arg Met Asn Tyr Val His  
 370 375 380  
 Arg Asp Leu Arg Ala Ala Asn Ile Leu Val Gly Glu Asn Leu Val Cys  
 385 390 395 400  
 Lys Val Ala Asp Phe Gly Leu Ala Arg Leu Ile Glu Asp Asn Glu Tyr  
 405 410 415  
 Thr Ala Arg Gln Gly Ala Lys Phe Pro Ile Lys Trp Thr Ala Pro Glu  
 420 425 430  
 Ala Ala Leu Tyr Gly Arg Phe Thr Ile Lys Ser Asp Val Trp Ser Phe  
 435 440 445  
 Gly Ile Leu Leu Thr Glu Leu Thr Thr Lys Gly Arg Val Pro Tyr Pro  
 450 455 460  
 Gly Met Val Asn Arg Glu Val Leu Asp Gln Val Glu Arg Gly Tyr Arg  
 465 470 475 480  
 Met Pro Cys Pro Pro Glu Cys Pro Glu Ser Leu His Asp Leu Met Cys  
 485 490 495  
 Gln Cys Trp Arg Arg Asp Pro Glu Glu Arg Pro Thr Phe Glu Tyr Leu  
 500 505 510  
 Gln Ala Phe Leu Glu Asp Tyr Phe Thr Ser Thr Glu Pro Gln Tyr Gln  
 515 520 525  
 Pro Gly Glu Asn Leu  
 530

<210> 4  
 <211> 2187  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> gene  
 <222> (1) .. (2187)

14/18

&lt;223&gt; human c-SRC cDNA

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (134)..(1483)

&lt;400&gt; 4

```

gcgccgcgctc ccgcaggccg tgatgccgcc cgcgcgaggagg tggcccggac cgcagtgtccc 60
caagagagagct ctaatggtac caagtgtacag gttggccttta ctgtgactcg gggacgccag 120
agctcctgag aag atg tca gca ata cag gcc gcc tgg cca tcc ggt aca 169
      Met Ser Ala Ile Gln Ala Ala Trp Pro Ser Gly Thr
              1              5              10

gaa tgt att gcc aag tac aac ttc cac ggc act gcc gag cag gac ctg 217
Glu Cys Ile Ala Lys Tyr Asn Phe His Gly Thr Ala Glu Gln Asp Leu
              15              20              25

ccc ttc tgc aaa gga gac gtg ctc acc att gtg gcc gtc acc aag gac 265
Pro Phe Cys Lys Gly Asp Val Leu Thr Ile Val Ala Val Thr Lys Asp
              30              35              40

ccc aac tgg tac aaa gcc aaa aac aag gtg ggc cgt gag ggc atc atc 313
Pro Asn Trp Tyr Lys Ala Lys Asn Lys Val Gly Arg Glu Gly Ile Ile
              45              50              55              60

cca gcc aac tac gtc cag aag cgg gag ggc gtg aag gcg ggt acc aaa 361
Pro Ala Asn Tyr Val Gln Lys Arg Glu Gly Val Lys Ala Gly Thr Lys
              65              70              75

ctc agc ctc atg cct tgg ttc cac ggc aag atc aca cgg gag cag gct 409
Leu Ser Leu Met Pro Trp Phe His Gly Lys Ile Thr Arg Glu Gln Ala
              80              85              90

gag cgg ctt ctg tac ccg ccg gag aca ggc ctg ttc ctg gtg cgg gag 457
Glu Arg Leu Leu Tyr Pro Pro Glu Thr Gly Leu Phe Leu Val Arg Glu
              95              100              105

agc acc aac tac ccc gga gac tac acg ctg tgc gtg agc tgc gac ggc 505
Ser Thr Asn Tyr Pro Gly Asp Tyr Thr Leu Cys Val Ser Cys Asp Gly
              110              115              120

aag gtg gag cac tac cgc atc atg tac cat gcc agc aag ctc agc atc 553
Lys Val Glu His Tyr Arg Ile Met Tyr His Ala Ser Lys Leu Ser Ile
              125              130              135              140

gac gag gag gtg tac ttt gag aac ctc atg cag ctg gtg gag cac tac 601
Asp Glu Glu Val Tyr Phe Glu Asn Leu Met Gln Leu Val Glu His Tyr
              145              150              155

acc tca gac gca gat gga ctc tgt acg cgc ctc att aaa cca aag gtc 649
Thr Ser Asp Ala Asp Gly Leu Cys Thr Arg Leu Ile Lys Pro Lys Val
              160              165              170

atg gag ggc aca gtg gcg gcc cag gat gag ttc tac cgc agc ggc tgg 697
Met Glu Gly Thr Val Ala Ala Gln Asp Glu Phe Tyr Arg Ser Gly Trp
              175              180              185

gcc ctg aac atg aag gag ctg aag ctg ctg cag acc atc ggg aag ggg 745
Ala Leu Asn Met Lys Glu Leu Lys Leu Leu Gln Thr Thr Ile Gly Lys Gly
              190              195              200

gag ttc gga gac gtg atg ctg ggc gat tac cga ggg aac aaa gtc gcc 793

```

15/18

Glu 205	Phe	Gly	Asp	Val	Met 210	Leu	Gly	Asp	Tyr	Arg 215	Gly	Asn	Lys	Val	Ala 220	
gtc	aag	tgc	att	aag	aac	gac	gcc	act	gcc	cag	gcc	ttc	ctg	gct	gaa	841
Val	Lys	Cys	Ile	Lys 225	Asn	Asp	Ala	Thr	Ala 230	Gln	Ala	Phe	Leu	Ala 235	Glu	
gcc	tca	gtc	atg	acg	caa	ctg	cgg	cat	agc	aac	ctg	gtg	cag	ctc	ctg	889
Ala	Ser	Val	Met	Thr 240	Gln	Leu	Arg	His 245	Ser	Asn	Leu	Val	Gln	Leu	Leu	
ggc	gtg	atc	gt	gag	gag	aag	ggc	ggg	ctc	tac	atc	gtc	act	gag	tac	937
Gly	Val	Ile 255	Val	Glu	Glu	Lys	Gly 260	Gly	Leu	Tyr	Ile	Val 265	Thr	Glu	Tyr	
atg	gcc	aag	ggg	agc	ctt	gtg	gac	tac	ctg	cgg	tct	agg	ggg	cgg	tca	985
Met	Ala 270	Lys	Gly	Ser	Leu	Val 275	Asp	Tyr	Leu	Arg	Ser 280	Arg	Gly	Arg	Ser	
gtg	ctg	ggc	gga	gac	tgt	ctc	ctc	aag	ttc	tcc	cta	gat	gtc	tgc	gag	1033
Val	Leu	Gly	Gly	Asp 290	Cys	Leu	Leu	Lys	Phe	Ser 295	Leu	Asp	Val	Cys	Glu 300	
gcc	atg	gaa	tac	ctg	gag	ggc	aac	aat	ttc	gtg	cat	cga	gac	ctg	gct	1081
Ala	Met	Glu	Tyr	Leu 305	Glu	Gly	Asn	Asn	Phe 310	Val	His	Arg	Asp	Leu 315	Ala	
gcc	cgc	aat	gtg	ctg	gtg	tct	gag	gac	aac	gtg	gcc	aag	gtc	agc	gac	1129
Ala	Arg	Asn	Val 320	Leu	Val	Ser	Glu	Asp 325	Asn	Val	Ala	Lys	Val 330	Ser	Asp	
ttt	ggt	ctc	acc	aag	gag	gcg	tcc	agc	acc	cag	gac	acg	ggc	aag	ctg	1177
Phe	Gly	Leu 335	Thr	Lys	Glu	Ala	Ser 340	Ser	Thr	Gln	Asp	Thr 345	Gly	Lys	Leu	
cca	gtc	aag	tgg	aca	gcc	cct	gag	gcc	ctg	aga	gag	aag	aaa	ttc	tcc	1225
Pro	Val	Lys	Trp	Thr	Ala 350	Pro 355	Glu	Ala	Leu	Arg	Glu 360	Lys	Lys	Phe	Ser	
act	aag	tct	gac	gtg	tgg	agt	ttc	gga	atc	ctt	ctc	tgg	gaa	atc	tac	1273
Thr	Lys	Ser	Asp	Val 370	Trp	Ser	Phe	Gly	Ile 375	Leu	Leu	Trp	Glu	Ile	Tyr 380	
tcc	ttt	ggg	cga	gtg	cct	tat	cca	aga	att	ccc	ctg	aag	gac	gtc	gtc	1321
Ser	Phe	Gly	Arg	Val 385	Pro	Tyr	Pro	Arg	Ile 390	Pro	Leu	Lys	Asp	Val 395	Val	
cct	cgg	gtg	gag	aag	ggc	tac	aag	atg	gat	gcc	ccc	gac	ggc	tgc	cgg	1369
Pro	Arg	Val 400	Glu	Lys	Gly	Tyr	Lys	Met 405	Asp	Ala	Pro	Asp	Gly 410	Cys	Pro	
ccc	gca	gtc	tat	gaa	gtc	atg	aag	aac	tgc	tgg	cac	ctg	gac	gcc	gcc	1417
Pro	Ala	Val 415	Tyr	Glu	Val	Met	Lys	Asn	Cys	Trp	His	Leu	Asp	Ala	Ala	
atg	cgg	ccc	tcc	ttc	cta	cag	ctc	cga	gag	cag	ctt	gag	cac	atc	aaa	1465
Met	Arg	Pro	Ser	Phe	Leu	Gln	Leu	Arg	Glu	Gln	Leu	Glu	His	Ile	Lys	
	430					435					440					
acc	cac	gag	ctg	cac	ctg	tgacggctgg	cctccgcctg	ggatcatgggc								1513
Thr	His	Glu	Leu	His	Leu 445											
					450											
ctgtggggac	tgaacctgga	agatcatgga	cctgggtgcc	ctgctcactg	ggccccgagcc											1573

16/18

tgaactgagc cccagcgggc tggcgggcct ttttcctgcg tcccagcctg caccctccg 1633  
 gccccgtctc tcttggaacc acctgtgggg cctggggagc ccactgaggg gccagggagg 1693  
 aaggaggcca cggagcggga ggcagcggcc caccacgtcg ggcttccctg gcctcccgcc 1753  
 actcgcttc ttagagtttt attcctttcc ttttttgaga ttttttttcc gtgtgtttat 1813  
 tttttattat ttttcaagat aaggagaaag aaagtaccca gcaaattgggc attttacaag 1873  
 aagtacgaat cttatttttc ctgtcctgcc cgtgaggggtg gggggggaccg ggccctctc 1933  
 tagggacccc tcgcccagc ctcattcccc attctgtgtc ccatgtcccg tgtctcctcg 1993  
 gtcgccccgt gtttgcgctt gaccatgttg cactgtttgc atgcgcccga ggcagacgtc 2053  
 tgtcaggggc ttggatttcg tgtgccgctg ccaccgccc acccgccctg tgagatggaa 2113  
 ttgtaataaa ccacgccatg aggacaccgc cgcccgccctc ggcgcttcc caccgaaaa 2173  
 aaaaaaaaaa aaaa 2187

<210> 5  
 <211> 450  
 <212> PRT  
 <213> Homo sapiens

<400> 5  
 Met Ser Ala Ile Gln Ala Ala Trp Pro Ser Gly Thr Glu Cys Ile Ala  
 1 5 10 15  
 Lys Tyr Asn Phe His Gly Thr Ala Glu Gln Asp Leu Pro Phe Cys Lys  
 20 25 30  
 Gly Asp Val Leu Thr Ile Val Ala Val Thr Lys Asp Pro Asn Trp Tyr  
 35 40 45  
 Lys Ala Lys Asn Lys Val Gly Arg Glu Gly Ile Ile Pro Ala Asn Tyr  
 50 55 60  
 Val Gln Lys Arg Glu Gly Val Lys Ala Gly Thr Lys Leu Ser Leu Met  
 65 70 75 80  
 Pro Trp Phe His Gly Lys Ile Thr Arg Glu Gln Ala Glu Arg Leu Leu  
 85 90 95  
 Tyr Pro Pro Glu Thr Gly Leu Phe Leu Val Arg Glu Ser Thr Asn Tyr  
 100 105 110  
 Pro Gly Asp Tyr Thr Leu Cys Val Ser Cys Asp Gly Lys Val Glu His  
 115 120 125  
 Tyr Arg Ile Met Tyr His Ala Ser Lys Leu Ser Ile Asp Glu Glu Val  
 130 135 140  
 Tyr Phe Glu Asn Leu Met Gln Leu Val Glu His Tyr Thr Ser Asp Ala  
 145 150 155 160  
 Asp Gly Leu Cys Thr Arg Leu Ile Lys Pro Lys Val Met Glu Gly Thr  
 165 170 175  
 Val Ala Ala Gln Asp Glu Phe Tyr Arg Ser Gly Trp Ala Leu Asn Met



[illegible]

```
<210> 6
<211> 14
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:9E10-myc
      epitope tag

<400> 6
```

18/18

Val Asp Met Glu Gln Lys Leu Ile Ala Glu Glu Asp Leu Asn  
1 5 10

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/11780

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : C12N 9/00

US CL : 435/183; 514/2

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/183; 514/2

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Extra Sheet.

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	MUKHOPADHYAY et al. Hypoxic induction of human vascular endothelial growth factor expression through c-Src activation. Nature. June 1995, Vol. 375, pages 577-581, especially abstract, page 578, col. 2, para. 2 through page 580, and Figs 3 and 4.	1-30 and 38
Y	KAPLAN et al. Association of the amino-terminal half of c-Src with focal adhesions alters their properties and is regulated by phosphorylation of tyrosine 527. EMBO J. 1994, Vol 13, pages 4745-4756, entire document	1-30, 37, 38

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*A* document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*B* earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family
*O* document referring to an oral disclosure, use, exhibition or other means	
*P* document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

Date of mailing of the international search report

18 OCT 1999

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231  
Facsimile No.Authorized officer  
Richard SchnitzerJOYCE BRIDGERS  
PARALEGAL SPECIALIST  
CHEMICAL MATRIS

Telephone No. (703) 308-0196



## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/11780

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	MUKHOPADHYAY et al. Wild-type p53 and v-Src exert opposing influences on human vascular endothelial growth factor gene expression. Cancer Research. 15 December 1995, Vol. 55, pages 6161-6165.	1-30, 38
Y	FLORIO et al. Aberrant protein phosphorylation at tyrosine is responsible for the growth inhibitory action of pp60v <sup>src</sup> expressed in the yeast saccharomyces cerevisiae. American Society For Cell Biology. March 1994, Vol. 5, pages 283-296, especially abstract	5-16, 21-30, 38
Y	US 5,264,618 A (FELGNER et al) 23 November 1993, col. 18, lines 18-33; and col. 20, lines 17-25.	14
Y	KOEGL et al. Generation of a temperature-sensitive cSRC. Virology. 1993, Vol. 196, pages 368-371, especially abstract.	1, 3, 12-16, and 37
Y	HIRAI et al. SH2 mutants of c-src that are host dependent for transformation are trans-dominant inhibitors of mouse cell transformation by activated c-src. Genes and Development. December 1990, Vol. 4, No. 12B, pages 2342-2352, abstract only.	1,3,12-16, 37

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/11780

## Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
  
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
  
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
  
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:  
1-30, 37, 38

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.  
☐ No protest accompanied the payment of additional search fees.

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US99/11780

### B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

#### MEDLINE

Src, K295M, Y527F, angiogenesis, (angiogen?(10w)(stimul? or generat? or caus? or improv? or therap?))

### BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be searched, the appropriate additional search fees must be paid.

Group I, claim(s) 1-30, 37 and 38, drawn to a src protein and a method of use.

Group II, claim(s) 1-16, drawn to a nucleic acid encoding a src protein.

Group III, claims 17-32, drawn to and methods of using nucleic acids encoding a src protein.

Group IV, claim 33, drawn to a viral gene transfer vector encoding a src protein.

Group V, claim 34, drawn to a non-viral gene transfer vector encoding a src protein.

Group VI, claim 35, drawn to a viral gene transfer vector encoding a src protein lacking kinase activity.

Group VII, claim 36, drawn to a non-viral gene transfer vector encoding a src protein lacking kinase activity.

The inventions listed as Groups I-VII do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons:

Claims 1-16 are drawn to a protein and a nucleic acid. PCT rule 13, 37 C.F.R. 1.475(c,d) does not allow for claims drawn to multiple independent products. If multiple products are claimed, the first invention of the category first mentioned in the claims, and the first recited invention of each of the other categories related to the first invention will be considered as the main invention. See PCT Article 17(3)(a) and 37 C.F.R. 1.476(c).

The technical feature of group I is the Src protein of claim 1. Group I is therefore composed of claims to the Src protein (claims 1-16, 37 and 38), and methods of using it (claims 17-30). The Src protein of claim 1, and genes encoding it, are well known in the art (see for example, Bricknell, Crit Rev Oncog 3: 401-446, 1992, abstract), therefore the claims have no special technical feature.

Groups II-VII are linked to group I because they comprise polynucleotides encoding the protein which is the technical feature of group I. However, because there can be no special technical feature, there is a lack of unity between the groups.